arXiv:1107.2707v1 [quant-ph] 14 Jul 2011

Structure of 2D Topological Stabilizer Codes

Héctor Bombín
Perimeter Institute for Theoretical Physics, 31 Caroline St. N., Waterloo, ON, N2L 2Y5, Canada

We provide a detailed study of the general structure of two-dimensional topological stabilizer quantum error correcting codes, including subsystem codes. Under the sole assumption of translational invariance, we show that all such codes can be understood in terms of the homology of string operators that carry a certain topological charge. In the case of subspace codes, we prove that two codes are equivalent under a suitable set of local transformations if and only they have equivalent topological charges. Our approach emphasizes local properties of the codes over global ones.

Contents

I. Introduction 1

II. Approach and results 2
A. Stabilizer codes 2
B. Topological codes and lattice Pauli groups 3
C. Independent generators and constraints 4
D. Charge 4
E. Strings and topological charge 5
F. Code structure and homology 6
G. Equivalence 7
H. Centralizers and 2D topology 7

III. Basic definitions 8
A. Notation 8
B. Lattice Pauli groups 10
C. Topological gauge groups 11
D. Examples 12

IV. Constraints and independent generators 12
A. Constraints 13
B. Independent generators 14

V. Charge 16
A. Charge groups 16
B. Charge of generators 16
C. Coarse graining 17
D. Strings 18
E. Charge and constraints 19

VI. Topological charge 20
A. String commutation rules 20
B. Invariance 23

VII. Topological Stabilizer Groups 23
A. Charge groups 23
B. Canonical charge generators 23
C. Characteristic of TSGs 24

VIII. Homological structure 24
A. Results 25
B. Examples 29

IX. Local equivalence of TSGs 30

X. Codes in the torus 31

XI. Conclusions and outlook 32

References 32

I. INTRODUCTION

Quantum error correcting codes [1–4] play a fundamental role in the quest to overcome the decoherence of quantum systems. Among them, stabilizer codes [5, 6] provide a large and flexible class of codes that are at the same time relatively easy to investigate. Interactions between physical qubits are sometimes subject to locality constraints in a geometrical sense. For example, qubits might be placed in a D-dimensional array with interactions available only between nearest neighbors. In such cases the subclass of topological stabilizer codes (TSC) [7–14] is a natural choice. TSCs do not only have very nice locality properties, but are also flexible in terms of the manipulation strategies that they allow.

The purpose of this paper is to investigate in full generality the structure of two-dimensional TSCs. In particular, not only subspace codes are considered but also the more general subsystem codes [15–17]. The only constraint imposed is translational symmetry of the bulk of the code, which provides a way to list the codes.

A stabilizer quantum error correcting code is described in terms of certain 'check operators', a set of commuting observables that have to be measured in order to get information about what errors have affected an encoded state. In TSCs these measurements are local in a geometrical sense. In particular, a TSC is in practice usually given as a recipe to construct the check operators on a given lattice of qubits. The lattice can be arbitrarily large, but check operators are defined locally, with support on a set of qubits contained in a bounded region. What makes these codes topological, as opposed to simply local, is that no information about the encoded qubits can be recovered if only access to a local set of qubits is granted. Indeed, operators on encoded qubits have support on a number of physical qubits that grows with the lattice size.

An essential feature of known topological codes is that they have an error threshold [8, 18–21]. That is, in the limit of large lattices, for noise below a certain threshold error correction is asymptotically perfect. This is true either in a simple error correction scenario or in a fault-tolerant scenario, and also when qubit losses are taken into account [22]. It is also offen true [23] —but for interesting exceptions, see[12, 14]— that the number of encoded qubits depends only on the homology of the qubit lattice. In particular a trivial homology gives no encoded qubits, but even in a planar setting it is possible to recover non-trivial codes by introducing suitable boundaries [24] and other defects such as twists [25]. Moreover, because in such 'homological' codes the lattices can be chosen very flexibly, it is possible to carry out certain computations by changing the code geometry over time, something called 'code deformation' [8, 26, 27]. Error correction turns out to be connected to classical statistical models [8, 11, 19] and this has given rise to fast algorithms to infer the errors [28]. Finally, there exist topological codes that are especially well suited to perform computations using transversal gates [9, 10], which minimizes error propagation.

General properties of TSCs in two or higher dimensions have already been explored to different degrees. General constraints on the code distance —directly related to the geometry of encoded operators— were first found in [29], and improved in [30]. The subsystem case was addresed in [31]. The geometry of logical operators in a subclass of TSCs, subject to constraints such as scale invariance of the number of encoded qubits, was studied in [23]. Constraints on the code distance for three-dimensional codes that do not satisfy this scale invariance condition have been recently developed [32].

One of the main results in this paper is that all two-dimensional TSCs can be understood in terms of 'string operators' and carry a 'topological charge'. In particular, homology plays an essential role and dictates the number of encoded qubits. In the case of subspace codes, the corresponding Hamiltonian model [7] —which has the code as its gapped ground state— exhibits anyonic excitations. Moreover, all subspace codes that give rise to the same anyon model turn out to be equivalent, in the sense that there exists a local transformation connecting them. The implications of this result, both from the condensed matter and quantum information perspectives, are explored in [33].

Our approach emphasizes local properties of the codes —such as the structure of check operators— over global ones —such as the number of encoded qubits—. In practice we realize this by considering infinite versions of the codes that cover the plane. The idea is that all operators acting on a finite number of qubits become automatically local in this infinite picture. At the same time, the homology of the plane is trivial and this simplifies the analysis.

This paper is divided as follows. Section II informally summarizes the main results, providing the intuition behind the main definitions and proofs. Section III introduces in a formal manner topological stabilizer groups (TSGs), used to model TSCs. Subsequent sections develop the results on TSGs, which are interpreted for TSCs in section X. Finally, section XI discusses natural extensions of the present work.

NOTE: To avoid repetition, we will most of the time omit the qualifier "two-dimensional" when discussing lattices, TSCs, and so on, but it should be understood in all cases.

II. APPROACH AND RESULTS

The purpose of this section is to summarize informally the approach taken to investigate TSCs and the results obtained. At the same time, it explains the motivation behind the main definitions and the intuition behind some proofs.

A. Stabilizer codes

Given a system with n qubits, its Pauli group is defined as

$$\mathcal{P}_n := \langle i\mathbf{1}, X_1, Z_1, \ldots, X_n, Z_n \rangle, \tag{1}$$

where X_i, Z_i denote the Pauli X and Z operators acting on the i-qubit. A stabilizer code [5, 6] on n qubits is defined by a subgroup of Pauli operators $\mathcal{S} \subset \mathcal{P}$, called stabilizer group, such that

$$-1 \notin \mathcal{S}. \tag{2}$$

Notice that $\mathcal{S}$ is then abelian and its elements are self-adjoint. The code is composed by those states $|\psi\rangle \in \mathbf{C}^{2n}$ such that

$$s|\psi\rangle = |\psi\rangle \tag{3}$$

for every stabilizer $s \in \mathcal{S}$. It is enough to check these conditions for a set $\{s_i\}$ of independent generators of $\mathcal{S}$. Such stabilizer generators serve as check operators: they can be measured in order to recover information about any errors affecting the encoded states. If $\mathcal{S}$ has $n-k$ independent generators the code subspace has dimension 2^k, it encodes k logical qubits. The logical or encoded Pauli operators are recovered as the quotient

$$\frac{\mathcal{Z}_{\mathcal{P}_n}(\mathcal{S})}{\mathcal{S}} \simeq \mathcal{P}_k, \tag{4}$$

where $\mathcal{Z}_{\mathcal{P}_n}(\mathcal{S})$ denotes the centralizer of $\mathcal{S}$ in $\mathcal{P}_n$. This centralizer is the group of undetectable Pauli errors: those that do not change the error syndrome. The quotient is necessary because the elements of $\mathcal{S}$ are trivial undetectable errors, without any effect on encoded states.

Given a set $\{s_i\}$ of generators of $\mathcal{S}$ we can write down a Hamiltonian

$$H = -\sum_i s_i. \tag{5}$$

Its ground subspace is the code subspace, and there is a gap of two energy units to any excited state. This gap is important when we consider families of TSCs with local generators in systems of arbitrary size, because the size independence of the gap gives rise to a gapped phase. Notice that an error syndrome in the code amounts to an excitation configuration in the Hamiltonian system.

It is also possible to encode only in a subsystem of the code subspace. In the stabilizer formalism this logical subsystem can be described as that in which the action of a certain gauge group $\mathcal{G} \subset \mathcal{P}$ is trivial [16]. The gauge group is such that

$$\mathcal{Z}_{\mathcal{G}}(\mathcal{G}) \propto \mathcal{S} \tag{6}$$

and we assume $\langle i\mathbf{1}\rangle \subset \mathcal{G}$. In this case logical operators are recovered as the quotient

$$\frac{\mathcal{Z}_{\mathcal{P}_n}(\mathcal{G})}{\mathcal{S}} \simeq \mathcal{P}_k, \tag{7}$$

where 2^k is no longer the dimension of the code subspace, which is instead 2^{k+r} for some r. Then $\mathcal{S}$ has $n-k-r$ independent generators and $\mathcal{G}$ has $n-k+r$ independent generators. Undetectable errors form the group $\mathcal{Z}_{\mathcal{P}_n}(\mathcal{S})$. They are, up to a product with an element of $\mathcal{G}$, logical operators. Indeed,

$$\frac{\mathcal{Z}_{\mathcal{P}_n}(\mathcal{G})}{\langle i\mathbf{1}\rangle\mathcal{S}} \simeq \frac{\mathcal{Z}_{\mathcal{P}_n}(\mathcal{S})}{\mathcal{G}}. \tag{8}$$

Stabilizer subsystem codes can be described just by giving $\mathcal{G}$, since then $\mathcal{S}$ is fixed up to signs that can be chosen arbitrarily.

For subsystem codes we no longer have such a straightforward interpretation in terms of a quantum Hamiltonian. A natural possibility is to take a set of generators $\{g_i\}$ of the gauge code as the Hamiltonian terms. This will guarantee that all the energy eigenvalues of the system have at least a degeneracy 2^k, since encoded operators commute with all Hamiltonian terms. However, there is no reason for an energy gap to persist in the limit of large system sizes.

B. Topological codes and lattice Pauli groups

For a TSC we mean a family of codes such that, loosely speaking,

- stabilizer generators are local, and
- non-trivial undetectable errors are non-local.

In addition, for subsystem codes we impose that

- gauge generators are local.

As explained in section X, certain topological subsystem codes naturally give rise to a few global generators either in the stabilizer or in the gauge group, but in such a way that it is irrelevant from the encoding perspective. Notice that there exist codes with local gauge generators and non-local stabilizer generators, such as Bacon-Shor codes [17]. We do not consider such codes topological, and indeed they do not have an essential feature of topological codes: a non-zero error threshold in the limit of large code size.

In order to characterize TSCs in a more concrete manner, let us first consider ordinary stabilizer codes, not subsystem ones. As discussed in the introduction, a TSC is usually given as a recipe to construct the check operators on a given lattice of qubits. We assume that this recipe, in the bulk of the code —e.g., ignoring possible boundaries or other defects— takes the form of a local and translationally invariant (LTI) set of check operators. Then we can consider an infinite version of the same lattice and construct using the recipe an infinite group with check operators as generators. We call this a lattice Pauli group (LPG). Notice that translational invariance is supposed to hold only at a suitably large coarse graining scale.

In general, for a LPG we mean a group that has as generators a LTI set of Pauli operators on a certain lattice of qubits. By definition, the Pauli group $\mathcal{P}$ on the infinite lattice of qubits is itself a LPG, with generators all the single qubit Pauli operators. The goal is to understand which properties distinguish those LPGs that correspond to a TSC, which we call topological stabilizer groups (TSG). Then we can shift our study of TSCs to that of TSGs.

We define TSGs as LPGs $\mathcal{S}$ satisfying (2) and the topological condition

$$\mathcal{Z}_{\mathcal{P}}(\mathcal{S}) \propto \mathcal{S}. \tag{9}$$

Since in the infinite lattice all the Pauli operators are local, this condition can be read:

Local undetectable errors do not affect encoded states.

To motivate (9) in detail, let $\mathcal{S}$ be the LPG corresponding to a certain TSC given by the family of stabilizer groups $\mathcal{S}_l$. Any operator $O \in \mathcal{Z}_{\mathcal{P}}(\mathcal{S})$ has an analog O' acting on the bulk of the l-th code in the family for l sufficiently large. Moreover, $O' \in \mathcal{Z}_{\mathcal{P}}(\mathcal{S}_l)$ due to the locality of the stabilizer generators, and no matter what detailed definition of locality we adopt, O' should be local for sufficiently large l. Since O' is undetectable and the codes $\mathcal{S}_l$ are topological, it follows that $\phi O' \in \mathcal{S}_l$ for some phase ϕ. Then if $\phi O \notin \mathcal{S}$ we can safely add ϕO, and all its translations, to an LTI generating set of $\mathcal{S}$, enlarging $\mathcal{S}$. The only question left then is whether one might have to keep adding larger and large generators, but this is not the case because $\mathcal{Z}_{\mathcal{P}}(\mathcal{S})$ has a LTI set of generators. This is guaranteed by corollary 13:

The centralizer of a LPG is a LPG.

Now consider the subsystem case. We define topological stabilizer subsystem groups (TSSGs) as LPGs $\mathcal{S}$

satisfying (2) and the topological condition

$$\mathcal{Z}_{\mathcal{P}}(\mathcal{Z}_{\mathcal{P}}(\mathcal{S})) \propto \mathcal{S}. \tag{10}$$

This condition is trivially satisfied when the number of qubits is finite, but is not generally true for LPGs, as an example in the next section shows. To motivate (10), let $\mathcal{S}$ be the LPG corresponding to a certain topological stabilizer subsystem code given by the family of stabilizer groups $\mathcal{S}_l$ and gauge groups $\mathcal{G}_l$. We first notice that it makes sense to define the gauge group as the LPG

$$\mathcal{G} := \mathcal{Z}_{\mathcal{P}}(\mathcal{S}). \tag{11}$$

Indeed, any operator $O \in \mathcal{Z}_{\mathcal{P}}(\mathcal{S})$ has a local analog $O' \in \mathcal{Z}_{\mathcal{P}}(\mathcal{S}_l)$ acting on the bulk of the l-th code in the family for l sufficiently large. Since O' is undetectable and the codes $(\mathcal{S}_l, \mathcal{G}_l)$ are topological, it follows that $\phi O' \in \mathcal{G}_l$ for some phase ϕ —but phases are unimportant in the gauge group—. Conversely, any operator O' of $\mathcal{G}_l$ with support in the bulk corresponds to an operator $O \in \mathcal{Z}_{\mathcal{P}}(\mathcal{S})$ due to the locality of the stabilizer generators. Second, due to the definition (11) we have $\mathcal{S} \subset \mathcal{G}$ and from that $\mathcal{Z}_{\mathcal{P}}(\mathcal{G}) \subset \mathcal{G}$ and $\mathcal{Z}_{\mathcal{P}}(\mathcal{G}) = \mathcal{Z}_{\mathcal{G}}(\mathcal{G})$. Any operator $O \in \mathcal{Z}_{\mathcal{G}}(\mathcal{G})$ has an analog $O' \in \mathcal{Z}_{\mathcal{G}_l}(\mathcal{G}_l)$ for l sufficiently large, and thus we can if needed enlarge $\mathcal{S}$ as above to get $\mathcal{S} \propto \mathcal{Z}_{\mathcal{P}}(\mathcal{G})$, which does not affect (11).

C. Independent generators and constraints

Given a LTI set G of generators of a LPG, a constraint is a subset of G such that the product of its elements is proportional to the identity —but none of the elements is—. We distinguish between local constraints, with a finite number of elements, and global constraints, with a possibly infinite number of elements. The latter are well defined thanks to locality. Elements of G are independent if they are not elements of a local constraint.

LPGs $\mathcal{A}$ that are the centralizer of some other LPG $\mathcal{B}$, so that $\mathcal{A} = \mathcal{Z}_{\mathcal{P}}(\mathcal{B})$, have very specific properties regarding independent generators and constraints, namely:

The centralizer of a LPG admits a LTI set of independent generators. Such a set of generators has a finite number of global constraints.

These results are lemma 15 and theorem 18. An intuitive explanation for the first one is provided in section II H.

For general LPGs it is not true that there exists always an LTI set of independent generators. As a counterexample —that also shows that (10) is not satisfied for any abelian LPG—, one can consider a square lattice with one qubit per site, and the LPG generated by nearest neighbor Pauli operators of the form $Z_i Z_j$. There is one generator per link, and they are not independent because the product of all the link operators in a closed circuit is the identity operator. Moreover, it is not difficult to see that it is not possible to choose a translationally invariant set of independent generators. Similarly, the result does not remain true for higher dimensions. A counterexample in dimension three is given by the 3D toric code [8], where the product of the plaquette operators forming a closed surface is the identity operator.

D. Charge

A central notion in this paper is that of charge. To introduce it, first consider the commutator of two Pauli elements

$$(p;q) := pqp^{-1}q^{-1} = \pm 1. \tag{12}$$

Given $p \in \mathcal{P}$ we can construct a group morphism

$$(p;\,\cdot\,) : \mathcal{P} \longrightarrow \pm 1. \tag{13}$$

Obviously $\langle i\mathbf{1}\rangle$ is a subset of the kernel of $(p;\,\cdot\,)$. So let us consider more generally, given a LPG $\mathcal{A}$, the set of morphisms $\Phi(\mathcal{A})$ from $\mathcal{A}$ to ± 1 that satisfy this property. They form an abelian group with product

$$(\phi\phi')(a) := \phi(a)\phi'(a), \qquad a \in \mathcal{A},\ \phi, \phi' \in \Phi(\mathcal{A}). \tag{14}$$

We are especially interested in the subgroup $\Phi^0(\mathcal{A}) \subset \Phi(\mathcal{A})$ of those morphisms such that the preimage of -1 contains a finite number of elements of any LTI set of generators. For example, if $\mathcal{A}$ is a TSG, then the elements of $\Phi^0(\mathcal{A})$ represent states of the corresponding Hamiltonian model with a finite number of excitations. On the other hand, if $\mathcal{A}$ is a TSSG we can no longer interpret the elements of $\Phi^0(\mathcal{A})$ as states, but just as abstractions of (13).

Since $(p;\,\cdot\,)(q;\,\cdot\,) = (pq;\,\cdot\,)$, the elements of $\Phi^0(\mathcal{A})$ of the form (13) form a subgroup, that we denote $\mathrm{Com}_{\mathcal{A}}(\mathcal{P})$. We are precisely interested in those elements of $\Phi^0(\mathcal{A})$ that do not correspond to a Pauli operator. It is natural to study them through the quotient

$$C_{\mathcal{A}} := \frac{\Phi^0(\mathcal{A})}{\mathrm{Com}_{\mathcal{A}}(\mathcal{P})}. \tag{15}$$

The original motivation for this definition comes from the case in which $\mathcal{A}$ is a TSG. In that case the equivalence classes can be regarded as classes of excitation configurations up to local transformations. That is, charge is conserved in any given region if the only transformations allowed are those that affect only that region. Interestingly, in the case of TSSGs it is useful to consider both the stabilizer charge group $C_{\mathcal{S}}$ and the gauge charge group $C_{\mathcal{G}}$.

For a centralizer LPG there exists a nice connection between charges and global constraints. Indeed, given a LTI set of generators we have

$$\#\ \textit{global constraints} \quad = \quad \#\ \textit{charges,}$$

as shown in corollary 26. In particular, the group of charges is finite. For TSSGs, $C_{\mathcal{S}}$ and $C_{\mathcal{G}}$ have the same number of elements

$$|C_{\mathcal{S}}| = |C_{\mathcal{G}}|. \tag{16}$$



FIG. 1: We represent sites as the faces of a square lattice. The striped sites form a path. Together with the grey sites they form the support of a corresponding string operator.



FIG. 2: Geometries involved in the definition of string commutation rules.

However, they are not related by a natural isomorphism. Instead, there is the natural morphism

$$C_\iota : C_\mathcal{G} \longrightarrow C_\mathcal{S} \tag{17}$$

derived from the restriction of morphisms from $\mathcal{G}$ to morphisms from $\mathcal{S}$. As we will see below, its kernel and the quotient of $C_\mathcal{S}$ with its image are related through a certain duality.

E. Strings and topological charge

Consider the charge group $C_\mathcal{G}$ of a TSSG. Given a LTI set of generators of $\mathcal{G}$, let us say that the support of $\phi \in \Phi(\mathcal{G})$ is the set of lattice sites for which there exist a generator g with support in the site and $\phi(g) = -1$. Since it is finite, we can coarse grain the lattice till each site can hold any of the charges, in the sense that there exists for any charge $c \in C_\mathcal{G}$ a morphism $\phi \in \Phi^0(\mathcal{G})$ with charge c and support at that site. Consider two such morphisms ϕ, ϕ' with the same charge c but support at different sites σ, σ'. Their product has trivial charge so that $\phi\phi' = (p; \cdot\,)$ for some $p \in \mathcal{P}$. Moreover, if we coarse grain enough we can choose p to have support along a string-shaped region —a thickened path, see Fig. 1— connecting σ and σ'. We say that p is a string operator with charge c and endpoints ϕ, ϕ'.

In the case of a TSG $\mathcal{S}$ and from the perspective of the Hamiltonian model, a string operator p of charge c transports a charge c from one endpoint to another. For example, when applied to a ground state p creates a pair of excitations with charge c on its endpoints. In the general case of a TSSG $\mathcal{S}$ we can no longer think of charges in terms of excitation configurations, but it is useful to visualize that a string operator transports a charge c.

Our next goal is to study the commutation properties of strings, which for many geometries only depend on their charges. We start considering three strings p, q, r with a shared endpoint and thus with a common charge c, see Fig. 2(a). The quantity

$$\theta(c) := (pq; pr) = (p;q)(p;r)(q;r) \tag{18}$$

only depends on the charge c, so that it is well defined. To check this, consider an alternative set of strings p', q', r' as in Fig. 2(a), together with the auxiliary strings u, v, w. Let $s = pqp'q'uv$ and $t = prp'r'uw$. Then $(s;t) = (pq;pr)(p'q';p'r')$, but $s, t \in \mathcal{Z}_\mathcal{P}(\mathcal{G}) \propto \mathcal{S}$ and thus $(s;t) = 1$ because $\mathcal{S}$ is abelian. We can interpret (18) as the topological spin of the charge c [34]. We say that a charge c is bosonic if $\theta(c) = 1$ and fermionic if $\theta(c) = -1$.

Now take two crossing strings p, q with charges c, c', as in Fig. 2(b). The mutual statistics of the charges c, d are given by the quantity [34]

$$\kappa(c,d) := (p;q). \tag{19}$$

It only depends on the charges c, d, so that it is well defined. The proof is analogous to the previous one, now with the geometry of Fig. 2(b). If $\kappa(c,d) = -1$ mutual statistics are said to be semionic, otherwise they are trivial. If in Fig. 2(c) p has charge c, q has charge d_1 and r has charge d_2, it follows that

$$\kappa(c, d_1 d_2) = \kappa(c, d_1)\kappa(c, d_2), \tag{20}$$

since $\kappa(c, d_1 d_2) = (p; qr) = (p;q)(p;r) = \kappa(c,d_1)\kappa(c,d_2)$.

The topological spin and mutual statistics are related. Consider the geometry of Fig. 2(d), where the strings p, q, r have charge c and the strings p', q', r' have charge d. The figure illustrates that

$$\theta(cd) = \theta(c)\theta(d)\kappa(c,d), \tag{21}$$

since $\theta(cd) = (pp'qq'; pp'rr') = (pq;pr)(p'q';p'r')(r;q') = \theta(c)\theta(d)\kappa(c,d)$. This gives in particular

$$\kappa(c,c) = 1. \tag{22}$$

In summary, the charges in $C_\mathcal{G}$ can be regarded as the topological charges of an anyon model, with fusion given by the charge group product. But the topological structure does not stop there, because it is possible to define κ for a string with charge in $C_{\mathcal{S}}$ and another with charge in $C_{\mathcal{G}}$. This works because $\mathcal{Z}_{\mathcal{P}}(\mathcal{S}) \subset \mathcal{Z}_{\mathcal{P}}(\mathcal{Z}_{\mathcal{P}}(\mathcal{G}))$, which is all we need for the construction in Fig. 2(b) to make sense. The morphism (17) preserves commutation, see proposition 32. Notice that we cannot define κ for two strings with charges in $C_{\mathcal{S}}$, nor can we define θ for charges in $C_{\mathcal{S}}$, unless of course we are dealing with a TSG rather than a general $TSSG$.

The fact that charges in $C_{\mathcal{G}}$ and $C_{\mathcal{S}}$ 'interact topologically' is crucial for the construction of canonical charge generators given in theorem 35, which we summarize next. First, there exist α, β such that

$$C_{\mathcal{G}} = \langle c_1, \dots, c_\alpha, d_1, \dots, d_\alpha, e_1, \dots, e_\beta \rangle, \tag{23}$$

where the e_i generate the kernel of C_ι, all mutual statistics between generators are trivial except for $\kappa(c_i, d_i) = -1$, and all generators are bosons except possibly e_β, c_α and d_α, always with $\theta(c_\alpha) = \theta(d_\alpha)$. Second, there exist $\tilde{e}_i$ such that

$$C_{\mathcal{S}} = \langle C_\iota(c_1), \dots, C_\iota(c_\alpha), C_\iota(d_1), \dots, C_\iota(d_\alpha), \tilde{e}_1, \dots, \tilde{e}_\beta \rangle \tag{24}$$

and we have the desired duality

$$\kappa(c_j, \tilde{e}_k) = \kappa(d_j, \tilde{e}_k) = 1, \quad \kappa(e_j, \tilde{e}_k) = 1 - 2\delta_{jk}. \tag{25}$$

Notice that the anyon model attached to $C_{\mathcal{G}}$ only has a few parameters, α, β, $\theta(c_\alpha)$ and $\theta(e_\beta)$. There exist TSSGs covering all possible combinations of these parameters, as shown in section VII B. In the case of TSGs $\beta = 0$ and thus we are left with the parameters α and $\theta(c_\alpha)$. The toric code has $\theta(c_\alpha) = 1$ and $\alpha = 1$, and thus combining N toric codes together we get a code with $\alpha = N$ and $\theta(c_\alpha) = 1$. On the other hand, no codes are known with $\theta(c_\alpha) = -1$. From the condensed matter perspective these would give rise to a chiral anyon model in a Hamiltonian with commuting terms, which is believed not to be possible [7].

F. Code structure and homology

With string operators and their geometrical commutation relations at hand, it is easy to uncover the general structure of TSSGs. Given a TSSG, we can put it in a standard form through the following process. The first step is to find a set of canonical charges $c_i, d_i, e_i, \tilde{e}_i$. Then we construct a translationally invariant framework of string operators, each in the form of a straight segment. For each charge there is a square lattice of segment operators, spatially disposed as indicated in Fig. 3. The idea is that the commutation relations between these segments operators are fixed by θ and κ —sometimes after suitable adjustments—.

To understand the structure of the code it is not necessary to visualize segment operators in the intricate framework of Fig. 3. Rather, it is enough to consider a single square lattice and its dual, see Fig. 4. Segment operators with charge c_i or e_i belong to the direct lattice, and those with charge d_i or $\tilde{e}_i$ to the dual lattice. We can form string operators by taking products of segment operators with the same charge, which thus can be regarded simply as $\mathbf{Z}_2$ chains of direct or dual segments.



FIG. 3: Framework of string segments. Each charge contributes a square lattice in which links are string operators with that charge.



FIG. 4: A square lattice and its dual. String operators can be visualized as collections of segments on these two lattices. We show for such strings: a direct plaquette, a dual plaquette, a direct boundary and a dual open string. The last two cross once.

The commutation properties of such string operators are simple. A string with charge $c \neq \tilde{e}_i$ anticommutes with another string of charge $c' \neq c$ if and only the two strings cross an odd number of times —see Fig. 4— and



FIG. 5: Logical operators corresponding to strings with charge c_i, in black, and strings with charge d_i, in grey, for any $i = 1, \dots, \alpha$. The topology is that of a torus, with opposite boundaries identified.

either $c = c_i$ and $c' = d_i$ (and viceversa) or $c = e_i$ and $c' = \tilde{e}_i$. Open string operators with charge e_i belong to $\mathcal{G}$. Closed strings operators —or boundary strings since the homology is trivial in the plane— with charge c_i, d_i, e_i belong to $\mathcal{S}$, and those with charge $\tilde{e}_i$ to $\mathcal{G}$. Any boundary operator is a product of 'plaquette operators', each of them the product of the segment operators forming a plaquette.

We take such elementary boundary operators as generators of $\mathcal{G}$ or $\mathcal{S}$. In particular, for $\mathcal{G}$ we take plaquette operators —see Fig. 4— constructed from strings with charge $c_i, d_i, \tilde{e}_i$, and for $\mathcal{S}$ we choose those with charge c_i, d_i, e_i. To form LTI sets of generators of $\mathcal{G}$ and $\mathcal{S}$ we then just need to add generators that have no charge content. In the case of $\mathcal{S}$ this means that these extra generators are unimportant in the global structure of the code, since their eigenvalue can be independently switched.

The homological perspective on string operators becomes most relevant when we go back to a code in a finite lattice. The simplest possible way to do this is by considering a finite lattice with periodic boundary conditions, so that the topology is that of a torus. This means that now closed string operators need not be boundaries, which has several consequences.

The first thing that we need to check is that the groups $\mathcal{G}$ and $\mathcal{S}$ generated by the corresponding local generators actually give rise to stabilizer subsystem code. As it turns out, when $\beta \neq 0$ a small adjustament is needed. We have to either add closed strings with charge e_i to $\mathcal{S}$ or closed string with charge $\tilde{e}_i$ to $\mathcal{G}$ —or a suitable mixture of these two options—. In practice the choice does not matter, and it certainly does not affect the number of encoded qubits, which depends only on α.

Indeed, each pair (c_i, d_i) contributes two logical qubits, so that $k = 2\alpha$. Logical operators for each pair can be chosen from homologically nontrivial string operators as in toric codes, see Fig. 5. String operators with the same charge and homologically equivalent have the same effect on encoded states, so that error correction can also be done as in toric codes [8], for each (c_i, d_i) pair separately [28] —although to achieve optimality correlations should be taken into account—. The syndrome (endpoints) for error string operators with charge c_i is given by plaquette operators that are strings with charge d_i, and viceversa.

G. Equivalence

An isomorphism between two TSGs is a translationally invariant morphism of the Pauli groups that preserves the elements of $\langle i\mathbf{1}\rangle$. Thus, we can also call it a LTI Clifford transformation, where the locality is a consequence of translational invariance. We say that two TSGs are equivalent if they can be related by a combination of isomorphisms, lattice coarse graining and addition/removal of disentangled qubits in a translationally invariant way—.

TSG isomorphisms preserve the topological charge structure, and the converse also holds, as shown in theorem 42:

Two TSGs are equivalent if and only if they have isomorphic topological charges.

This result is a consequence of the general structure of TSGs discussed above.

H. Centralizers and 2D topology

To finish this summary of results, let us explain the role played by the topology of the plane and the fact that the LPG is a centralizer for the existence of an LTI set of independent generators. Let the support of a constraint be the union of the support of its elements. Consider a LPG $\mathcal{A}$ of the form $\mathcal{A} = \mathcal{Z}_{\mathcal{P}}(\mathcal{B})$. In order to recover a LTI set of independent generators from an arbitrary LTI set of generators of $\mathcal{A}$, all we need to do is remove in a translationally invariant way a sufficient amout of generators. If a generator is part of a constraint, it can be removed. The removal of constrained generators with support in the regions A, marked with dark grey in Fig. 6(a), can be done without much difficulty. It is enough to choose a large enough separation between these regions, so that generators in different regions are subject to independent constraints, which allows a translationally invariant removal. After the removal it is unclear whether the generators with support in different regions B, marked with light grey in Fig. 6(a), are also subject to independent constraints. But given a constraint R with support for example in the region B_0, marked with stripes in Fig. 6(a), we can find —as shown below— another constraint R_0 containing the same generators as R in B_0



(a)

(b)

(c)

(d)

FIG. 6: The regions involved in the construction of an LTI set of independent generators of a LPG that is the centralizer of a LPG, see the text.

but with no support in the rest of regions B. This shows that generators with support in different regions B are indeed subject to independent constraints. Then we are left with the regions C, marked with white in Fig. 6(a), but these are disconnected from each other and we can deal with them as follows. The constraints that are left can only include generators with support in regions C now, and since they are distant enough and the generators are local, the restriction of any constraint to a particular region C is still a constraint. Thus, generators with support in different regions C are also subject to independent constraints.

Thus, the crucial point is the existence, for any given constraint R with support in regions B and C, of a constraint R_0 with the same support as R in B_0 but no support in other regions B. To guarantee it, we take the size of the regions A as large as needed, so that near B_0 any such R has support only in a long and narrow band. Then, up to translations, the elements in R must repeat along the band at certain stretches of a length larger than the size of the generators. At each side of the band we pick two such repeated stretches, marked with stripes in Fig. 6(b), and form a new constraint R_∞ that extendes to infinity by repeating the set of constraints in between the repeated stretches up to translations. This is depicted in Fig. 6(c), where the places where the repeated stretches overlap are marked with stripes and the support of R_∞ is darkened. We can then consider a subset of R_∞, containing only the elements on the darkened region in Fig. 6(d), which produces a set of generators S with product $p_1p_2 \in \mathcal{P}$, where p_i have each support in one of the two regions marked with stripes. Since $p_1p_2 \in \mathcal{Z}_\mathcal{P}(\mathcal{B})$ and $\mathcal{B}$ has local generators, it follows that $p_1, p_2 \in \mathcal{Z}_\mathcal{P}(\mathcal{B})$ since the supports of p_1 and p_2 are sufficiently far appart. There exist sets of generators S_i of $\mathcal{A}$, $i = 1, 2$, with supports in a small neighborhood of the support of p_i and such that their product is proportional to p_i. Then to obtain R_0 it is enough to combine S, S_1 and S_2 in a suitable way —namely, through symmetric difference—.

III. BASIC DEFINITIONS

This section intends to put together most of the constructions needed later, with the aim of both presenting them and serving as a reference for later sections. It ends with a list of basic examples of topological codes.

A. Notation

Sets — For integers we use the notation $\mathbf{N}_0 := \{0\} \cup \mathbf{N}^*$. $|S|$ denotes the cardinality of a set S, $\mathbf{P}(S)$ its power set and $\mathbf{P}_0(S) := \{s \in \mathbf{P}(S) \,:\, |s| < \aleph_0\}$. Given a mapping $f\,:\, A \longrightarrow B$ and any subsets $a \subset A$, $b \subset B$, we denote by $f[a] \subset B$ the image of a and by $f^{-1}[b] \subset A$ the preimage of b. Given any set S, we regard $\mathbf{P}(S)$ as the abelian group with addition given by the symmetric difference

$$S_1 + S_2 := S_1 \cup S_2 - S_1 \cap S_2, \qquad S_1, S_2 \subset S. \tag{26}$$

2D lattice — Since we are interested in topological aspects, we consider without loss of generality square lattices. Sites are labeled as usual with integer coordinates $(j,k) \in \mathbf{Z}^2$, with $\Sigma = \mathbf{Z}^2$ the set of all sites, and $\mathbf{0} := (0,0)$. A block is a set of sites of the form

$$\square_{m,n}^{m',n'} := \{(j,k) \,:\, m \leq j < m'; n \leq k < n'\}. \tag{27}$$

Given a set of sites $\gamma \in \mathbf{P}_0(\Sigma)$, $||\gamma||$ denotes the smallest $k \in \mathbf{N}_0$ such that $\gamma \subset \square_{m,n}^{m+k,n+k}$ for some $m,n \in \mathbf{Z}$. For any $\gamma \subset \Sigma$ and $l \in \mathbf{N}_0$ we define a 'thickened' set

$$\mathrm{Thk}^l(\gamma) := \bigcup_{(i,j)\in\gamma} \square_{i-l,j-l}^{i+l+1,j+l+1}, \tag{28}$$

and set $\mathrm{Thk}(\gamma) := \mathrm{Thk}^1(\gamma)$.

Connectedness — Two sites (i,j) and (m,n) are adjacent if $|m-i| \leq 1$ and $|n-j| \leq 1$. An ordered list of sites $(\sigma_k)_{k=1}^n = (\sigma_0, \cdots, \sigma_n)$ where σ_k is adjacent to σ_{k+1} for all $k = 1, \ldots, n$ is a path from σ_0 to σ_n. We define the inverse γ^{-1} of a path γ in the usual way, an also the composition of paths $\gamma' \circ \gamma$, where the last site of γ is the first site of γ'. A set of sites $\gamma \subset \Sigma$ is connected if for any $\sigma, \sigma' \in \gamma$ there exists a path $(\sigma_k)_{k=1}^n$ with $\sigma_0 = \sigma$, $\sigma_n = \sigma'$ and $\sigma_i \in \gamma$, $i = 1, \ldots, n$. Given an ordered set of sites $(\sigma_k)_{k=1}^n = ((a_k, b_k))_{k=1}^n$, let $\sigma'_k := (a_{k+1}, b_k)$, $k = 1, \ldots, n-1$ and define

$$\mathrm{Path}(\sigma_1, \ldots, \sigma_n) \tag{29}$$

to be the path starting at σ_1, moving in a straight line to σ'_1, then in a straight line to σ_2, then $\sigma'_2, \ldots$, and finally ending at σ_n. When no confusion is possible we abuse notation and identify a path (σ_i) with the set of sites $\{\sigma_i\}$.

Homology — Let Γ be an infinite square lattice. Γ_{fc}, Γ^*_{fc}, Γ_{edg} and Γ^*_{edg} denote, respectively, the set of faces, dual faces, edges and dual edges. We represent $\mathbf{Z}_2$ chains of these objects as elements of $\mathbf{P}(\Gamma_{\text{fc}})$, $\mathbf{P}(\Gamma^*_{\text{fc}})$, $\mathbf{P}(\Gamma_{\text{edg}})$ and $\mathbf{P}(\Gamma^*_{\text{edg}})$. ∂ denotes the usual boundary operation, which we express in particular as

$$\begin{aligned} \mathbf{P}(\Gamma_{\text{fc}}) &\xrightarrow{\partial} \mathbf{P}(\Gamma_{\text{edg}}) \xrightarrow{\partial} \mathbf{P}(\Gamma^*_{\text{fc}}), \\ \mathbf{P}(\Gamma^*_{\text{fc}}) &\xrightarrow{\partial} \mathbf{P}(\Gamma^*_{\text{edg}}) \xrightarrow{\partial} \mathbf{P}(\Gamma_{\text{fc}}). \end{aligned} \tag{30}$$

Define the sets of direct and dual cycles (or cycles and co-cycles)

$$\begin{aligned} \Gamma_{\text{cyc}} &:= \{E \subset \Gamma_{\text{edg}} \,:\, \partial E = \emptyset\}, \\ \Gamma^*_{\text{cyc}} &:= \{E \subset \Gamma_{\text{edg}} \,:\, \partial E = \emptyset\}. \end{aligned} \tag{31}$$

Given an edge e, it dual is denoted e^*. We name the elements $\mathbf{e}_1, \mathbf{e}_2 \in \Gamma_{\text{edg}}$, $\mathbf{f} \in \Gamma_{\text{fc}}, \mathbf{f}^* \in \Gamma^*_{\text{fc}}$ according to Fig. 7. In the case of faces, the symbol * is just a label.

We will also consider finite and periodic square lattices Γ, thus with the topology of a torus. In that case we choose certain particular homologically nontrivial cycles $\mathbf{z}_1, \mathbf{z}_2 \in \Gamma_{\text{cyc}}$, $\mathbf{z}^*_1, \mathbf{z}^*_2 \in \Gamma^*_{\text{cyc}}$. In particular, $\mathbf{z}_1$ and $\mathbf{z}^*_2$ wind once horizontally, and $\mathbf{z}_2$ and $\mathbf{z}^*_1$ wind once horizontally.



FIG. 7: Naming some elements of Γ.

Qubits and operators — Sites hold a common and finite number of qubits. We denote by $X^\alpha_{j,k}$, $Z^\alpha_{j,k}$ the single qubit X and Z operators acting on the qubit with label α on the site (j,k). They generate for each site α an algebra of operators A_α. Consider the infinite tensor product $A := \otimes_\alpha A_\alpha$, with elements $\otimes_\alpha a_\alpha$ with all but a finite number of the product terms identical $a_\alpha = \mathbf{1}$. The Pauli group $\mathcal{P} \subset A$ is

$$\mathcal{P} := \langle i\mathbf{1}\rangle\langle X^\alpha_{ij}, Z^\alpha_{ij}\rangle_{i,j,\alpha}. \tag{32}$$

Given a Pauli operator $p \in \mathcal{P}$ we denote by $\mathrm{Sup}(p)$ the set of sites containing qubits on the support of p, that is, on which p acts nontrivially. The range of p is

$$||p|| := ||\mathrm{Sup}(p)||. \tag{33}$$

For $\gamma \subset \Sigma$ we let $p|_\gamma$ be the restriction of p to qubits in γ, defined only up to a global phase. Given $p, q \in \mathcal{P}$ we denote the commutator by

$$(p;q) := pqp^{-1}q^{-1} = \pm 1. \tag{34}$$

Given subgroups $\mathcal{A}, \mathcal{B} \subset \mathcal{P}$, the centralizer of $\mathcal{A}$ in $\mathcal{B}$ is

$$\mathcal{Z}_{\mathcal{B}}(\mathcal{A}) := \{b \in \mathcal{B} \,:\, \forall a \in \mathcal{A} \quad (a;b) = 1\}. \tag{35}$$

We write $\mathcal{A} \propto \mathcal{B}$ meaning $\langle i\mathbf{1}\rangle\mathcal{A} = \langle i\mathbf{1}\rangle\mathcal{B}$.

Pauli group subsets — For $S \subset \mathcal{P}$, $\gamma \subset \Sigma$ and $q \in \mathcal{P}$ we define

$$\begin{aligned} \mathrm{Sup}(S) &:= \bigcup \mathrm{Sup}[S], \\ S|_\gamma &:= \{s \in S \,:\, \mathrm{Sup}(s) \subset \gamma\}, \\ S||_\gamma &:= \{s \in S \,:\, \mathrm{Sup}(s) \cap \gamma \neq \emptyset\}, \\ (S;q) &:= \prod_{p \in S_q} (p;q), \end{aligned} \tag{36}$$

where $S_q = \{p \in S\,:\, \mathrm{Sup}(p)\cap\mathrm{Sup}(q)\neq\emptyset\}$. The range of S is

$$||S|| := ||\mathrm{Sup}(S)||. \tag{37}$$

We define a group morphism

$$\begin{aligned}\mathrm{Pro}\,:\; \mathbf{P}_0(\mathcal{P}) &\longrightarrow \mathcal{P}/\langle i\mathbf{1}\rangle\\ S &\longmapsto \langle i\mathbf{1}\rangle \prod_{p\in S} p\end{aligned} \tag{38}$$

where the ordering of the terms in the product is immaterial. We say that a set of generators $\mathcal{A}_g$ of a group $\mathcal{A}\subset\mathcal{P}$ is independent if the kernel of the restriction of this morphism to $\mathbf{P}_0(\mathcal{A}_g)$ is a subgroup of $\mathbf{P}(\langle i\mathbf{1}\rangle)$.

Morphisms — Let $\mathcal{A}\subset\mathcal{P}$ be a group. The group morphisms

$$\phi\,:\,\mathcal{A}\longrightarrow\{1,-1\} \tag{39}$$

such that $\phi(i)=1$ form an abelian group $\Phi(\mathcal{A})$ setting

$$(\phi\phi')(a) := \phi(a)\phi'(a), \qquad a\in\mathcal{A},\ \phi,\phi'\in\Phi(\mathcal{A}). \tag{40}$$

Given a subset $\mathcal{A}_g$ of $\mathcal{A}$, we define a group morphism

$$\begin{aligned}\overline{\mathrm{Ker}_{\mathcal{A}_g}}\;:\;\Phi(\mathcal{A}) &\longrightarrow \mathbf{P}(\mathcal{A}_g - \langle i\mathbf{1}\rangle)\\ \phi &\longmapsto \mathcal{A}_g\cap\phi^{-1}[-1],\end{aligned} \tag{41}$$

mapping ϕ to the complement of its kernel in $\mathcal{A}_g$. If $\mathcal{A}_g$ is an independent set of generators of $\mathcal{A}$ this is an isomorphism. We denote the inverse by

$$\mathrm{Mor}_{\mathcal{A}_g}\,:\,\mathbf{P}(\mathcal{A}_g-\langle i\mathbf{1}\rangle)\longrightarrow\Phi(\mathcal{A}). \tag{42}$$

The support of $\phi\in\Phi(\mathcal{A})$ with respect to $\mathcal{A}_g$ is

$$\mathrm{Sup}_{\mathcal{A}_g}(\phi) := \mathrm{Sup}(\overline{\mathrm{Ker}_{\mathcal{A}_g}}(\phi)). \tag{43}$$

We define the subgroup of morphisms

$$\Phi^0(\mathcal{A}_g) := \{\phi\in\Phi(\langle\mathcal{A}_g\rangle)\,:\,|\mathrm{Sup}_{\mathcal{A}_g}(\phi)|<\aleph_0\}. \tag{44}$$

Given $\phi\in\Phi^0(\mathcal{A}_g)$, we will denote with the same name the group morphism

$$\begin{aligned}\phi\;:\;\mathbf{P}(\mathcal{A}_g) &\longrightarrow \{1,-1\}\\ S &\longmapsto (-1)^{|S\cap\overline{\mathrm{Ker}_{\mathcal{A}_g}}(\phi)|}\end{aligned} \tag{45}$$

The commutation structure gives rise to the group morphisms

$$\begin{aligned}\mathrm{Com}_{\mathcal{A}}\,:\,\mathcal{P}&\longrightarrow\Phi(\mathcal{A}) & \mathrm{Com}_{\mathcal{A}}\,:\,\mathbf{P}(\mathcal{P})&\longrightarrow\Phi(\mathcal{A})\\ p&\longmapsto(p;\cdot) & S&\longmapsto(S;\cdot)\end{aligned} \tag{46}$$

For simplicity, for $p\in\mathcal{P}$ we set

$$\mathrm{Sup}_{\mathcal{A}_g}(p) := \mathrm{Sup}_{\mathcal{A}_g}(\mathrm{Com}_{\langle\mathcal{A}_g\rangle}(p)). \tag{47}$$

Translation operators — We define translation operators for sites setting

$$T_{m,n}(j,k) := (j+m, k+n). \tag{48}$$

The action on edges or faces of Γ is analogous. For Pauli operators we define

$$T_{m,n}\,:\,\mathcal{P}\longrightarrow\mathcal{P} \tag{49}$$

as the Pauli group automorphisms such that

$$T_{m,n}(i\mathbf{1}) := i\mathbf{1}, \tag{50}$$

$$T_{m,n}(X^\alpha_{j,k}) := X^\alpha_{j+m,k+n}, \tag{51}$$

$$T_{m,n}(Z^\alpha_{j,k}) := Z^\alpha_{j+m,k+n}. \tag{52}$$

Finally, for $\phi\in\Phi(\mathcal{A})$ we set

$$(T_{m,n}(\phi))(\cdot) := \phi(T_{-m,-n}(\cdot)). \tag{53}$$

To generalize translation to sets of translatable objects, we recursively define whenever it makes sense

$$T_{m,n}(A) := \{T_{m,n}(a)\,:\,a\in A\}. \tag{54}$$

Finally, we set for $d\in\mathbf{N}^*$

$$T^{(d)}(\cdot) := \bigcup_{m,n\in\mathbf{Z}} T_{md,nd}(\cdot). \tag{55}$$

Coarse graining and composition — A qubit lattice can be coarse grained by identifying each block $\square^{(m+1)l,(n+1)l}_{ml,nl}$ as the site (m,n) of the new lattice, for some $l\in\mathbf{N}^*$. We denote the effect of coarse graining on a given object x by $\mathrm{Crs}_l(x)$, be it an operator, a set of sites, a group or any other structure.

Two disjoint qubit lattices can be put together to form a single lattice. If $\mathcal{A}$ is a Pauli subgroup on the first lattice and $\mathcal{B}$ is a Pauli subgroup on the second lattice we define a new group

$$\mathcal{A}\otimes\mathcal{B} := \{a\otimes b\,:\,a\in\mathcal{A}, b\in\mathcal{B}\}. \tag{56}$$

When discussing lattice composition, for simplicity we write p instead of $\mathbf{1}\otimes p$ or $p\otimes\mathbf{1}$.

B. Lattice Pauli groups

In what follows, consider a 2D square lattice of qubits with Pauli group (32).

Definition 1 *Let S be a set such that* $\mathrm{Sup}[S]$ *and* $T_{m,n}[S]$ *are defined. S is local and translationally invariant (LTI) if there exist $k,l\in\mathbf{N}^*$ such that for any $s\in S$, $m,n\in\mathbf{Z}$,*

$$||s||\leq k, \qquad T_{ml,nl}(s)\in S. \tag{57}$$

In that case we say that S is k-bounded and has step l.

Definition 2 *A subgroup $\mathcal{A} \subset \mathcal{P}$ is a lattice Pauli group (LPG) if it admits a LTI set of generators $\mathcal{A}_g \subset \mathcal{A}$,*

$$\mathcal{A} = \langle \mathcal{A}_g \rangle. \tag{58}$$

When we say that $\mathcal{A}_g$ generates a LPG $\mathcal{A}$ it is understood that $\mathcal{A}_g$ is a LTI set.

The following result is an immediate consequence of the definition of LPGs.

Proposition 3 *Consider a set of LPGs $\bar{\mathcal{A}}^1, \dots, \bar{\mathcal{A}}^n$ on a given lattice. There exists $L \in \mathbf{N}^*$ such that the coarse grained LPGs*

$$\mathcal{A}^k = \mathrm{Crs}_L(\bar{\mathcal{A}}^k) \subset \mathcal{P}, \qquad k = 1, \dots, n, \tag{59}$$

admit LTI sets of independent generators $\mathcal{A}_g^k$ such that

1. *$\mathcal{A}_g^i$ has step 1, and*
2. *$||a|| \leq 2$ for any $a \in \mathcal{A}_g^i$.*

We will assume that all the LPGs that we discuss satisfy these properties, unless otherwise stated.

Definition 4 *Given two LPGs $\mathcal{A}, \mathcal{B}$ defined on disjoint qubit lattices, their composition is the LPG*

$$\mathcal{A} \otimes \mathcal{B} \tag{60}$$

defined on the union of the two lattices.

Notice that $\mathcal{A} \otimes \mathcal{B}$ is indeed an LPG because $\mathcal{A}_g \cup \mathcal{B}_g$ is a valid LTI generating set.

Definition 5 *A LPG isomorphism from $\mathcal{A}_1 \subset \mathcal{P}_1$ to $\mathcal{A}_2 \subset \mathcal{P}_2$ is a group isomorphism*

$$F : \mathcal{P}_1 \longrightarrow \mathcal{P}_2 \tag{61}$$

such that

$$F[\mathcal{A}_1] = \mathcal{A}_2, \qquad F(i) = i, \tag{62}$$

and, for any $m, n \in \mathbf{Z}$ and some $d \in \mathbf{N}^*$,

$$T_{md,nd} \circ F = F \circ T_{md,nd}\,. \tag{63}$$

We say that F has step d and denote it as an arrow

$$\mathcal{A}_1 \xrightarrow{F} \mathcal{A}_2. \tag{64}$$

The range $||F||$ of F is

$$||F|| := \max_{\sigma \in \square_{0,0}^{d,d}} ||\{\sigma\} \cup \mathrm{Sup}(F[\mathcal{P}_1|_{\{\sigma\}}])||. \tag{65}$$

Two LPGs $\mathcal{A}_1$ and $\mathcal{A}_2$ are isomorphic if there exists such an isomorphism F. Notice that $||F||$ is finite and such that

$$\mathrm{Sup}(F(p)) \subset \mathrm{Thk}^{||F||-1}(\mathrm{Sup}(p)) \tag{66}$$

for every $p \in \mathcal{P}_1$. Also, F preserves commutators

$$(\,\cdot\,;\,\cdot\,) = (F(\,\cdot\,); F(\,\cdot\,)). \tag{67}$$

Given the LPG isomorphism (64) we can coarse grain both lattices, that of $\mathcal{A}$ and that of $\mathcal{B}$, and obtain an isomorphism

$$\mathrm{Crs}_l(\mathcal{A}) \xrightarrow{\mathrm{Crs}_l(F)} \mathrm{Crs}_l(\mathcal{B}), \tag{68}$$

where $l \in \mathbf{N}^*$. Given two LPG isomorphisms

$$\mathcal{A}_1 \xrightarrow{F} \mathcal{A}_2, \qquad \mathcal{B}_1 \xrightarrow{G} \mathcal{B}_2, \tag{69}$$

if the corresponding lattices are disjoint we form in the obvious way the isomorphism

$$\mathcal{A}_1 \otimes \mathcal{B}_1 \xrightarrow{F \otimes G} \mathcal{A}_2 \otimes \mathcal{B}_2. \tag{70}$$

Thus we can consider the strict monoidal category **LPG** of lattice Pauli groups. A trivial example of a functor over this category is the coarse graining operator. But our main object of study is a much more interesting functor over a closely related category. In the tradition of algebraic topology, this functor removes unnecessary information in such a way that isomorphic elements have the same image. We will even be able to show that for a given subcategory, elements are isomorphic iff they have the same image under the functor. Although we will sometimes use the language of category theory, we do not use it as a tool or go into details.

C. Topological gauge groups

Definition 6 *A lattice stabilizer group is a LPG $\mathcal{S} \subset \mathcal{P}$ such that*

$$-1 \notin \mathcal{S}. \tag{71}$$

Definition 7 *A topological stabilizer group (TSG) is a lattice stabilizer group $\mathcal{S} \subset \mathcal{P}$ such that*

$$\mathcal{Z}_{\mathcal{P}}(\mathcal{S}) \propto \mathcal{S}. \tag{72}$$

Definition 8 *A topological stabilizer subsystem group (TSSG) is a lattice stabilizer group $\mathcal{S} \subset \mathcal{P}$ such that*

$$\mathcal{Z}_{\mathcal{P}}(\mathcal{Z}_{\mathcal{P}}(\mathcal{S})) \propto \mathcal{S}. \tag{73}$$

Its gauge group is

$$\mathcal{G} := \mathcal{Z}_{\mathcal{P}}(\mathcal{S}). \tag{74}$$

As we will show in corollary 13, $\mathcal{G}$ is a LPG. Notice that TSGs are TSSGs. The product $\mathcal{S}_1 \otimes \mathcal{S}_2$ of two TSSGs is also a TSSG. A TSSG isomorphism F from $\mathcal{S}_1$ to $\mathcal{S}_2$ is just a LPG isomorphism. Then F is also a LPG isomorphism from $\mathcal{G}_1$ to $\mathcal{G}_2$.



(a) (b) (c)

FIG. 8: Unit cells for the TPGs in the examples. (a) In the toric code, qubits can be identified with horizontal and vertical links in a square lattice, or with vertical and horizontal links in its dual. Stabilizers are related to faces and dual faces. (b) In the honeycomb subsystem code qubits can be identified with the vertices of a honeycomb lattice. Gauge generators are related to links. (c) In the topological subsystem color code qubits can be identified with the vertices of a lattice derived from the honeycomb. The numbering is obtained by moving counterclockwise along any hexagonal plaquette. Gauge generators are related to links.

D. Examples

Empty code — This is the most simple TSG, with no qubits.

Trivial code — This TSG is constructed on a lattice with one qubit per site. We denote by $\mathcal{P}_{\mathrm{T}}$ the corresponding pauli group. The stabilizer $\mathcal{S}_{\mathrm{T}}$ has a set of independent generators

$$\mathcal{S}_g^{\mathrm{T}} := T^{(1)}\left(\{Z_{0,0}\}\right). \tag{75}$$

Subsystem trivial code — This TSSG is constructed on the same lattice as the trivial code. The stabilizer is trivial, $\mathcal{S}_{\mathrm{S}} = \{\mathbf{1}\}$ and the gauge group is $\mathcal{G}_{\mathrm{S}} = \mathcal{P}_{\mathrm{T}}$.

Toric code — This TSG [7] is constructed on a lattice with two qubits per site, labeled as h and v. We denote by $\mathcal{P}_{\mathrm{TC}}$ the corresponding pauli group. The stabilizer $\mathcal{S}_{\mathrm{TC}}$ has a set of independent generators

$$\mathcal{S}_g^{\mathrm{TC}} := T^{(1)}\left(\{S^X, S^Z\}\right), \tag{76}$$

where

$$\begin{aligned} S^X &:= X^h_{0,0} X^h_{-1,0} X^v_{0,0} X^v_{0,-1}, \\ S^Z &:= Z^h_{0,0} Z^h_{0,1} Z^v_{0,0} Z^v_{1,0}. \end{aligned} \tag{77}$$

Subsystem toric code — This TSSG is constructed on the same lattice as the toric code. The stabilizer $\mathcal{S}_{\mathrm{STC}}$ has a set of independent generators

$$\mathcal{S}^{\mathrm{STC}} := T^{(1)}\left(\{S^Z\}\right). \tag{78}$$

The gauge group $\mathcal{G}_{\mathrm{STC}}$ has a set of independent generators

$$\mathcal{G}_g^{\mathrm{STC}} := \langle i\mathbf{1}\rangle T^{(1)}\left(\{S^X, Z^h_{0,0}, Z^v_{0,0}\}\right). \tag{79}$$

Honeycomb subsystem code — This TSSG [13] is constructed on a lattice with two qubits per site, labeled as 1,2. The stabilizer $\mathcal{S}_{\hexagon}$ has a set of independent generators

$$\mathcal{S}_g^{\hexagon} := T^{(1)}\left(\{S^{\hexagon}\}\right), \tag{80}$$

where

$$S^{\hexagon} := X^2_{0,0} Z^1_{1,0} Y^2_{1,0} X^1_{1,1} Y^1_{0,1} Z^2_{0,1}. \tag{81}$$

The gauge group $\mathcal{G}_{\hexagon}$ has a set of independent generators

$$\mathcal{G}_g^{\hexagon} := \langle i\mathbf{1}\rangle T^{(1)}\left(\{G^X, G^Y, G^Z\}\right), \tag{82}$$

where

$$\begin{aligned} G^X &:= X^1_{0,0} X^2_{0,0}, \\ G^Y &:= Y^1_{1,0} Y^2_{0,0}, \\ G^Z &:= Z^1_{0,1} Z^2_{0,0}, \end{aligned} \tag{83}$$

Topological subsystem color code — This TSSG [11] is constructed on a lattice with six qubits per site, labeled 1-6. The stabilizer $\mathcal{S}_{\mathrm{col}}$ has a set of independent generators

$$\mathcal{S}_g^{\mathrm{col}} := T^{(1)}\left(\{S^1, S^2\}\right), \tag{84}$$

where

$$\begin{aligned} S^1 &:= Z^1_{0,0} Z^2_{1,0} Z^3_{1,0} Z^4_{1,1} Z^5_{0,1} Z^6_{0,1}, \\ S^2 &:= X^1_{0,0} Y^2_{1,0} X^3_{1,0} Y^4_{1,1} X^5_{0,1} Y^6_{0,1} \cdot \\ &\quad \cdot Y^1_{0,1} Y^2_{0,1} Y^3_{0,0} Y^4_{1,0} Y^5_{1,0} Y^6_{1,1} \cdot \\ &\quad \cdot X^1_{1,0} X^2_{1,1} X^3_{0,1} X^4_{0,1} X^5_{0,0} X^6_{1,0}. \end{aligned} \tag{85}$$

The gauge group $\mathcal{G}_{\mathrm{col}}$ has a set of independent generators

$$\mathcal{G}_g^{\mathrm{col}} := \langle i\mathbf{1}\rangle T^{(1)}\left(\{G^1, G^2, \dots, G^{10}\}\right), \tag{86}$$

where

$$\begin{aligned} G^1 &:= X^1_{0,0} Y^2_{1,0}, & G^2 &:= X^2_{1,0} Y^3_{1,0}, \\ G^3 &:= X^3_{1,0} Y^4_{1,1}, & G^4 &:= X^4_{1,1} Y^5_{0,1}, \\ G^5 &:= X^5_{0,1} Y^6_{0,1}, & G^6 &:= X^6_{0,1} Y^1_{0,0}, \\ G^7 &:= Z^1_{0,0} Z^5_{0,0}, & G^8 &:= Z^3_{0,0} Z^5_{0,0}, \\ G^9 &:= Z^2_{0,0} Z^4_{0,0}, & G^{10} &:= Z^2_{0,0} Z^6_{0,0}. \end{aligned} \tag{87}$$

IV. CONSTRAINTS AND INDEPENDENT GENERATORS

The ultimate goal of this section is to show how a LTI set of independent generators can be constructed for any LPG that is the centralizer of a LPG. $\mathcal{A} = \langle \mathcal{A}_g \rangle$, $\mathcal{B} = \langle \mathcal{B}_g \rangle$ denote LPGs on a given 2D lattice. $\mathcal{P}_g$ denotes the set of single qubit Pauli operators.



FIG. 9: Illustration of proposition 10. The region γ_l appears in darker grey. The closed shapes represents the support of elements of $\overline{\mathrm{Ker}_{\mathcal{B}_g}}(\phi)$. The support of S is contained in the larger rectangle.

A. Constraints

Definition 9 *The groups of $\mathcal{B}$-constraints of $\mathcal{A}_g$ are*

$$\begin{aligned}\mathrm{Cnstr}_{\mathcal{B}}(\mathcal{A}_g) &:= \{S \subset \mathcal{A}_g - \langle i\mathbf{1}\rangle \,:\, \mathrm{Com}_{\mathcal{B}}(S) = 1\},\\ \mathrm{Cnstr}^0_{\mathcal{B}}(\mathcal{A}_g) &:= \mathbf{P}_0(\mathcal{A}_g) \cap \mathrm{Cnstr}_{\mathcal{B}}(\mathcal{A}_g)\end{aligned} \tag{88}$$

We use the notation

$$\begin{aligned}\mathrm{Cnstr}(\mathcal{A}_g) &:= \mathrm{Cnstr}_{\mathcal{P}}(\mathcal{A}_g),\\ \mathrm{Cnstr}^0(\mathcal{A}_g) &:= \mathrm{Cnstr}^0_{\mathcal{P}}(\mathcal{A}_g),\end{aligned} \tag{89}$$

so that

$$\begin{aligned}\mathrm{Cnstr}(\mathcal{A}_g) &= \{S \subset \mathcal{A}_g - \langle i\mathbf{1}\rangle \,:\, (S;\cdot\,) = 1\},\\ \mathrm{Cnstr}^0(\mathcal{A}_g) &= \{S \in \mathbf{P}_0(\mathcal{A}_g - \langle i\mathbf{1}\rangle) \,:\, \mathrm{Pro}(S) = \langle i\mathbf{1}\rangle\}.\end{aligned} \tag{90}$$

Notice that $\mathcal{A}_g$ is independent if and only if

$$\mathrm{Cnstr}^0(\mathcal{A}_g) = \{\emptyset\}. \tag{91}$$

Proposition 10 *For any $m \in \mathbf{N}^*$ there exists $n \in \mathbf{N}_0$ such that for every $\phi \in \mathrm{Com}_{\mathcal{B}}[\mathcal{A}]$ and $l \in \mathbf{N}_0$ with*

$$\mathrm{Sup}_{\mathcal{B}_g}(\phi) \subset \mathcal{B}_g||_{\gamma_l}, \qquad \gamma_l := \square^{l,m}_{0,0}, \tag{92}$$

there exists $S \in \mathbf{P}_0(\mathcal{A}_g)$ with $\phi = \mathrm{Com}_{\mathcal{B}}(S)$ and

$$\mathrm{Sup}(S) \subset \mathrm{Thk}^n(\gamma_l). \tag{93}$$

We denote by $N(\mathcal{A}_g, \mathcal{B}_g, m) \in \mathbf{N}_0$ the smallest of such n, see Fig. 9.

Proof — Let $\mathcal{B}_1 := \langle \mathcal{B}_g||_{\gamma_1}\rangle$. For $\phi \in \Phi(\mathcal{B}_1)$ let

$$A_\phi := \{a \in \mathcal{A} \,:\, \mathrm{Sup}_{\mathcal{B}_g}(a) \subset \mathcal{B}_g||_{\gamma_\infty} \,\wedge\, \mathrm{Com}_{\mathcal{B}_1}(a) = \phi\} \tag{94}$$

and $\Phi_1 := \{\phi \in \Phi_{\mathcal{B}_1} \,:\, A_\phi \neq \emptyset\}$. Φ_1 is finite because $\mathcal{B}_1$ is finite. For each $\phi \in \Phi_1$ we choose $a_\phi \in A_\phi$ that satisfies $\mathrm{Sup}_{\mathcal{B}_g}(a_\phi) \subset \gamma_l$ for a minimal l and let l_p be this minimum value. We also choose $S_\phi \in \mathbf{P}_0(\mathcal{A}_g)$ with $\mathrm{Com}_{\mathcal{B}}(a_\phi) = \mathrm{Com}_{\mathcal{B}}(S_\phi)$. Finally, we take the minimal $n \in \mathbf{N}_0$ such that for any $\phi \in \Phi_1$

$$\mathrm{Sup}(S_\phi) \subset \mathrm{Thk}^n(\gamma_{l_p}). \tag{95}$$

Take any $a \in \mathcal{A}$ with $\mathrm{Sup}_{\mathcal{B}_g}(a) \subset \mathcal{B}_g||_{\gamma_l}$ and let $\phi = \mathrm{Com}_{\mathcal{B}_1}(a)$. Then $a = a_\phi a'$ with $\mathrm{Sup}_{\mathcal{B}_g}(a') \subset \mathcal{B}_g||_{T_{1,0}\,(\gamma_{l-1})}$. The result follows by induction on l. ☐

Corolary 11 *For any $m \in \mathbf{N}^*$ there exists $n \in \mathbf{N}_0$ such that for every $a \in \mathcal{A}$ with*

$$\mathrm{Sup}(a) \subset \gamma_l := \square^{m,l}_{0,0}, \tag{96}$$

there exists $S \in \mathbf{P}_0(\mathcal{A}_g)$ with $a \in \mathrm{Pro}(S)$ and

$$\mathrm{Sup}(S) \subset \mathrm{Thk}^n(\gamma_l). \tag{97}$$

We denote by $N(\mathcal{A}_g, m) \in \mathbf{N}_0$ the smallest of such n.

Proof — Notice that $\mathrm{Com}_{\mathcal{P}}[\mathcal{A}] \simeq \mathcal{A}$ and $\mathrm{Sup}_{\mathcal{P}_g}(\mathrm{Com}_{\mathcal{P}}(\cdot)) = \mathrm{Sup}(\cdot)$. ☐

Lemma 12 *$\mathrm{Cnstr}^0_{\mathcal{B}}(\mathcal{A}_g)$ admits a LTI set of generators.*

Proof — Let $n_1 = N(\mathcal{A}_g, \mathcal{B}_g, 2)$ according to proposition 10 and $n_2 = N(\mathcal{A}_g, \mathcal{B}_g, 2)$ according to the axis-exchanged version of proposition 10. Set $N = \max(n_1, n_2)$. We claim that

$$G := \{R \in \mathrm{Cnstr}^0_{\mathcal{B}}(\mathcal{A}_g) \,:\, ||R|| \leq L\}, \tag{98}$$

where $L := 8N + 2$, generates $\mathrm{Cnstr}^0_{\mathcal{B}}(\mathcal{A}_g)$.

Take any $R \in \mathrm{Cnstr}^0_{\mathcal{B}}(\mathcal{A}_g)$. If $||R|| \leq L$ then $R \in \langle G\rangle$. In other case, without lost of generality we assume $\mathrm{Sup}(R) \subset \square^{a,b}_{0,0}$ with $a > b$ and $a > L$. Let $l = 4N + 10$ and set $S = S_1 + S_2$ with $S_1 = S|_{\square^{l,b}_{0,0}}$. Then $\mathrm{Sup}(S_2) \subset \square^{a,b}_{l-2,0}$, $\mathrm{Com}_{\mathcal{B}}(S_1) = \mathrm{Com}_{\mathcal{B}}(S_2) =: \phi \in \mathrm{Com}_{\mathcal{B}}[\mathcal{A}]$, and

$$\mathrm{Sup}_{\mathcal{B}_g}(\phi) \subset \mathrm{Sup}_{\mathcal{B}_g}(S_1) \cap \mathrm{Sup}_{\mathcal{B}_g}(S_2) \subset \mathcal{B}_g||_{\square^{l,b}_{l-2,0}}. \tag{99}$$

Due to proposition 10 and translational symmetry, there exists $S_3 \in \mathbf{P}_0(\mathcal{A}_g)$ with $\phi = \mathrm{Com}_{\mathcal{B}}(S_3)$ and $\mathrm{Sup}(S_3) \subset \mathrm{Thk}^N(\square^{l,b}_{l-2,0})$. Then $S = S_1' + S_2'$, where $S_1' = S_1 + S_3$ and $S_2' = S_2 + S_3$ are elements of $\mathrm{Cnstr}^0_{\mathcal{B}}(\mathcal{A}_g)$. But $S_i' \in \square^{a_i,b_i}_{u_i+a_i,v_i+b_i}$, $i = 1, 2$, with $a_1 = l + N$, $b_1 = b + 2N$, $a_2 = a + 1 + N$ and $b_2 = b + 2N$. Thus $a_i + b_i < a + b$ and by induction on $a + b$ it follows that $R \in \langle G\rangle$. ☐

Corolary 13 *$\mathcal{Z}_{\mathcal{A}}(\mathcal{B})$ is a LPG.*

Proof — Notice that $\mathcal{Z}_{\mathcal{A}}(\mathcal{B}) \propto \bigcup \mathrm{Pro}[\mathrm{Cnstr}^0_{\mathcal{B}}(\mathcal{A})]$. ☐

Proposition 14 *Let $\mathcal{A} = \mathcal{Z}_{\mathcal{P}}(\mathcal{B})$. For any $a \in \mathcal{A}$ and $\gamma \in \mathbf{P}_0(\Sigma)$*

$$\mathrm{Sup}(a) \cap (\mathrm{Thk}(\gamma) - \gamma) = \emptyset \quad\Longrightarrow\quad a|_\gamma \in \langle i\mathbf{1}\rangle\mathcal{A}. \tag{100}$$

Proof — Since $\mathcal{B}_g$ is 2-bounded, for any $b \in \mathcal{B}_g$ with $\mathrm{Sup}(b) \cap \gamma \neq \emptyset$ we have $\mathrm{Sup}(b) \cap (\mathrm{Sup}(a) - \gamma) = \emptyset$ and thus $(b; a|_\gamma) = (b; a) = 1$. $\Box$

Lemma 15 *Let $\mathcal{A} = \mathcal{Z}_\mathcal{P}(\mathcal{B})$ and $\mathcal{A}_g$ be independent.* $\mathrm{Cnstr}(\mathcal{A}_g)$ *is finite.*

Proof — Let $n_1 := N(\mathcal{A}_g, 2)$ according to corollary 11 and $n_2 := N(\mathcal{A}_g, 2)$ according to the axis-exchanged version of corollary 11. Set $N = \max(n_1, n_2)$ and $D = 2N + 2$. First, we want to show that for any $R \in \mathrm{Cnstr}(\mathcal{A}_g)$ we have

$$R||_{\Box_{0,0}^{2,D}} = \emptyset \quad \Rightarrow \quad R||_{\Box_{2,N}^{\infty,D-N}} = \emptyset. \tag{101}$$

We start observing that for any $l \in \mathbf{N}^* - \{1\}$, there exist $k_1, k_2 \in \mathbf{N}^*$ with $k_2 - 2 > k_1 > l$ and such that

$$\left(T_{k_2-k_1,0}\,(R) + R\right)||_{\Box_{k_2,0}^{k_2+2,D}} = \emptyset, \tag{102}$$

because the set

$$\{S \subset \mathcal{A}_g \,:\, S = S||_{\Box_{0,0}^{2,D}}\} \tag{103}$$

is finite. Let

$$R_0 := R||_{\Box_{2,0}^{\infty,D}}, \qquad R_1 := T_{k_2-k_1,0}\,(R_0) + R_0. \tag{104}$$

Clearly we have

$$R_1||_{\Box_{0,0}^{2,D} \cup \Box_{k_2,0}^{k_2+2,D}} = \emptyset. \tag{105}$$

Then $R_2 := R_1||_{\Box_{2,0}^{k_2,D}}$ satisfies $\mathrm{Sup}(\mathrm{Pro}(R_2)) \subset \gamma_1 \cap \gamma_2$ for

$$\gamma_1 := \Box_{0,-2}^{k_2+2,0}, \quad \gamma_2 := \Box_{0,D}^{k_2+2,D+2}. \tag{106}$$

According to proposition 14 there exist $a_i \in \mathcal{A}$, $i = 1, 2$, such that $a_1 a_2 \in \mathrm{Pro}(R_2)$. Applying corollary 11, there exist $S_i \in \mathbf{P}_0(\mathcal{A}_g)$, $i = 1, 2$ with $a_i \in \mathrm{Pro}(S_i)$ and $\mathrm{Sup}(S_i) \subset \mathrm{Thk}^N(\gamma_i)$. Then $R_3 = R_2 + S_1 + S_2 \in \mathrm{Cnstr}^0(\mathcal{A}_g) = \{\emptyset\}$ and $R_1||_{\Box_{2,N}^{l,D-N}} = R_3||_{\Box_{2,N}^{l,D-N}} = \emptyset$. This can only be true if $R||_{\Box_{2,N}^{l,D-N}} = \emptyset$.

We next show that for any $R_1, R_2 \in \mathrm{Cnstr}(\mathcal{A}_g)$ with

$$R_1||_{\Box_{0,0}^{2,D}} = R_2||_{\Box_{0,0}^{2,D}}, \tag{107}$$

we have $R_1 = R_2$, which implies the finiteness of $\mathrm{Cnstr}(\mathcal{A}_g)$ because (103) is finite. Let $R = R_1 + R_2$. Then according to the above result $R||_{\Box_{2,N}^{\infty,D-N}} = \emptyset$. Moreover, we can apply the result with the first coordinate inverted and a suitable translation, getting $R||_{\Box_{-\infty,N}^{\infty,D-N}} = \emptyset$. But now for any $j \in \mathbf{Z}$ we have $R||_{\Box_{j,N}^{j+D,2+N}} = \emptyset$. Applying the same reasoning with the coordinates exchanged gives $R||_{\Box_{j+N,-\infty}^{j+D-N,\infty}} = \emptyset$ for any $j \in \mathbf{Z}$. Thus $R_1 + R_2 = \emptyset$. $\Box$



FIG. 10: Illustration of proposition 16. R has no support on the region γ, in medium grey. The support of S is contained in the light grey area, which extends to infinity on both sides. S and R coincide in the region γ_{-n}^n, in dark grey.

B. Independent generators

Proposition 16 *Let $\mathcal{A}_g$ be M-bounded. For any $m, n \in \mathbf{N}_0$ there exists $l \in \mathbf{N}_0$ such that the following holds. For any $R \in \mathrm{Cnstr}(\mathcal{A}_g)$ satisfying*

$$R||_\gamma = \emptyset, \qquad \gamma := \Box_{-l,-M}^{l,0} \cup \Box_{-l,m}^{l,m+M}, \tag{108}$$

there exists $S \in \mathrm{Cnstr}(\mathcal{A}_g)$ such that

$$S \subset \mathcal{A}_g|_{\gamma_{-\infty}^\infty}, \quad (S + R)||_{\gamma_{-n}^n} = \emptyset, \quad \gamma_a^b := \Box_{a,0}^{b,m}. \tag{109}$$

We denote by $L(\mathcal{A}_g, m, n)$ the minimal such l, see Fig. 10

Proof — For $a < b$, we set $\gamma_b^a := \gamma_a^b$.

Let $l \in \mathbf{N}_0$ be such that for any $R \in \mathrm{Cnstr}(\mathcal{A}_g)$ satisfying (108) there exist $m_x, n_x \in \mathbf{N}_0$, $x = \pm$, with $n \leq x m_x < x m_x + M \leq x n_x \leq r - M$ and such that

$$\left(T_{d_x,0}\,(R) + R\right)||_{\gamma_{n_x}^{n_x + xM}} = \emptyset, \quad d_x := n_x - m_x. \tag{110}$$

There exists such l because the set $\mathcal{A}_g||_{\gamma_0^M}$ is finite and $\mathcal{A}_g||_{\gamma_t^M} = T_{t,0}\,(\mathcal{A}_g||_{\gamma_0^M})$. We claim that this choice for l satisfies the statement in the lemma. To check it, given R, m_x, n_x as above let

$$\begin{aligned} R_x &:= R||_{\gamma_{m_x}^{n_x}} - R||_{\gamma_{m_x - xM}^{m_x}}, \\ R_x^\infty &:= \bigcup_{a \in \mathbf{N}^*} T_{ad_x,0}\,(R_x), \\ S &:= R||_{\gamma_{n_-}^{n_+}} \cup R_-^\infty \cup R_+^\infty. \end{aligned} \tag{111}$$

Due to (108), $R||_{\gamma_{n_-}^{n_+}} \subset R|_{\mathrm{Thk}^M(\gamma_{n_-}^{n_+})} \subset R|_{\gamma \cup \gamma_{-r}^r} \subset \mathcal{A}_g|_{\gamma_{-\infty}^\infty}$ and thus $S \subset \mathcal{A}_g|_{\gamma_{-\infty}^\infty}$. The definition of S ensures that

$$(S + R)||_{\gamma_{n_- - M}^{n_+ + M}} = \emptyset, \tag{112}$$

and thus in particular $(S + R)||_{\gamma_{-n}^n} = \emptyset$. The definition is also such that for any $a \in \mathbf{N}^*$, $x = \pm$,

$$(T_{ad_x,0}\,(S) + S)||_{\gamma_{m_x + ad_x}^\infty} = \emptyset \tag{113}$$

To complete the proof, it suffices to show that for any $p \in \mathcal{P}_g$ we have $(S;p)=1$. Let $\mathrm{Sup}(p) \subset \Box_{t,-\infty}^{t+1,\infty}$ for some $t \in \mathbf{N}_0$. W.l.o.g. we set $t \geq 0$. If $t \leq n_+$, $(S;p)=(R;p)=1$ according to (112). In other case $t = m_+ + ad_+ + t'$ for some $a \in \mathbf{N}^*$, $0 \leq t' < d_+$. Using (113) we get $(S;p) = (T_{ad_+,0}\,(S);p) = (S;T_{-ad_+,0}\,(p)) = 1$. $\Box$

Proposition 17 *Let $\mathcal{A}_g$ be a LTI set of step d and $\mathrm{Cnstr}^0(\mathcal{A}_g) = \langle R_0 \rangle$ for some LTI set $R_0 \subset \mathbf{P}_0(\mathcal{A}_g)$. Let $\gamma \in \mathbf{P}_0(\Sigma)$ be such that for any $S \in R_0$ there is at most a pair $(m,n) \in \mathbf{Z}^2$ such that*

$$S||_{T_{md,nd}\,(\gamma)} \neq \emptyset. \tag{114}$$

Then there exists $A \subset \mathcal{A}_g||_\gamma$ such that

$$\mathcal{A} = \langle \mathcal{A}'_g \rangle, \qquad \mathcal{A}'_g := \mathcal{A}_g - T^{(d)}\,(A), \tag{115}$$

and $S||_{T^{(d)}\,(\gamma)} = \emptyset$ for any constraint $S \in \mathrm{Cnstr}^0(\mathcal{A}'_g)$.

Proof — If $S||_\gamma = \emptyset$ for any constraint $S \in \mathrm{Cnstr}^0(\mathcal{A}_g)$, clearly $A = \emptyset$ suffices. In other case, choose $a \in \mathcal{A}_g||_\gamma$ and $S \in R_0$ such that $a \in S$. We claim that $\mathcal{A} = \langle \mathcal{A}'_g \rangle$ with $\mathcal{A}'_g := \mathcal{A}_g - T^{(d)}\,(a)$. Due to translational invariance, it suffices to show $a \in \langle \mathcal{A}'_g \rangle$. But $a \in \mathrm{Pro}(S - \{a\})$, and since $T^{(d)}\,(a) \cap S = \{a\}$ we have $(S - \{a\}) \subset \mathcal{A}'_g$.

Let $S_{m,n} := T_{md,nd}\,(S)$. Given $S' \in R_0$, let $f(S') = S' + S_{m,n}$ if $T_{md,nd}\,(a) \in S'$, $f(S') = S'$ otherwise. Then the set $R'_0 = f[R_0]$ generates $\mathrm{Cnstr}^0(\mathcal{A}'_0)$. Moreover, $\mathcal{A}'_g$ and R'_0 satisfy the conditions of the lemma, because $\mathrm{Sup}(S + S_{m,n}) \subset \mathrm{Sup}(S) \cup \mathrm{Sup}(S_{m,n})$. Since $|(\mathcal{A}'_g||_\gamma)| = |(\mathcal{A}_g||_\gamma)| - 1$, the result follows by induction on $|(\mathcal{A}_g||_\gamma)|$. $\Box$



FIG. 11: The regions involved in the proof of theorem 18. One of them, γ_2^X is only outlined with a dashed contour.

Theorem 18 *Let $\mathcal{A} = \mathcal{Z}_{\mathcal{P}}(\mathcal{B})$. Then $\mathcal{A}$ admits a LTI set of independent generators.*

Proof — We will construct from $\mathcal{A}_g$ a LTI set of independent generators $\mathcal{A}_3$ through a series of intermediate sets $\mathcal{A}_i$ so that

$$\mathcal{A}_3 \subset \mathcal{A}_2 \subset \mathcal{A}_1 \subset \mathcal{A}_0 := \mathcal{A}_g \tag{116}$$

At each step we will apply proposition 17 to a given region γ_i and obtain

$$\mathcal{A}_{i+1} = \mathcal{A}_i - T^{(D)}\,(A_i) \tag{117}$$

for a suitable $D \in \mathbf{N}^*$, $A_i \subset \mathcal{A}_i||_{\gamma_i}$. Let us define $R_i := \mathrm{Cnstr}^0(\mathcal{A}_i)$. Since we apply proposition 17, we have $S||_{T^{(D)}\,(\gamma_i)} = \emptyset$ for any constraint $S \in R_j$ if $j > i$. The goal is thus to have

$$\{\emptyset\} = R_3 \subset R_2 \subset R_1 \subset R_0. \tag{118}$$

R_g^i will denote a LTI set of generators of $R_i$of step D.

0) W.l.o.g. we assume that $\mathcal{A}_g^0$ and R_g^0 have step 1 and are 2-bounded. Let $n_1 := N(\mathcal{A}_g, 2)$ according to corollary 11 and $n_2 := N(\mathcal{A}_g, 2)$ according to the axis-exchanged version of corollary 11. Set $N = \max(1, n_1, n_2)$ and $K := 2N+2$. Let $l_1 = L(\mathcal{A}_g, K, 2)$ according to proposition 16 and $l_2 = L(\mathcal{A}_g, K, 2)$ according to the axis-exchanged version of proposition 16. Set $L := \max(l_1, l_2)$ and $D := K + 2L + 2N$. The geometry of the proof, see Fig. 11, involves the regions

$$\begin{aligned}
\gamma_1^X &:= \Box_{K+L-2,0}^{K+L+2,K}, & \gamma_1^Y &:= \Box_{0,K+L-2}^{K,K+L+2},\\
\gamma_2^X &:= \Box_{N,0}^{D+K-N,K}, & \gamma_2^Y &:= \Box_{0,N}^{K,D+K-N},\\
\gamma_3^X &:= \Box_{N,0}^{K-N,K}, & \gamma_3^Y &:= \Box_{0,N}^{K,K-N},\\
\gamma_4^X &:= T_{D,0}\,(\gamma_3^X), & \gamma_4^Y &:= T_{D,0}\,(\gamma_3^X),\\
\gamma_5^X &:= \mathrm{Thk}^N(\gamma_3^X \cup \gamma_4^X), & \gamma_5^Y &:= \mathrm{Thk}^N(\gamma_3^Y \cup \gamma_4^Y),\\
\gamma_6^X &:= \gamma_2^X \cup \gamma_5^X, & \gamma_6^Y &:= \gamma_2^Y \cup \gamma_5^Y,\\
\gamma_0 &:= \Box_{K,K}^{D,D}, & \gamma_1 &:= \gamma_1^X \cup \gamma_1^Y,
\end{aligned}$$

$$\gamma_2 := \Box_{2-L,0}^{K+L-2,K} \cup \Box_{0,2-L}^{K,K+L-2}. \tag{119}$$

1) $\mathcal{A}_0$, R_g^0 and γ_0 satisfy the conditions of proposition 17 and we get $\mathcal{A}_1$ setting $i=0$ in (117). Let for $\alpha = X, Y$

$$Q^\alpha := \{S \in R_1 \,:\, S||_{\gamma_1^\alpha} \neq \emptyset \;\wedge\; S|_{\gamma_6^\alpha} \neq S\}. \tag{120}$$

We claim that $R_1 = \langle R_g^1 \rangle$ with

$$R_g^1 := R^1 - T^{(D)}\,(Q^X \cup Q^Y). \tag{121}$$

Take any $R \in R_1$. The set

$$\{\gamma \in T^{(D)}\,(\{\gamma_1^X, \gamma_1^Y\}) \,:\, R||_\gamma \neq \emptyset\} \tag{122}$$

is finite, and thus we proceed by induction on its cardinality. If it is empty $R \in R_1$. In other case, due to translation and axis-exchange symmetries we can assume that $S||_{\gamma_1^X} \neq \emptyset$. Then it suffices to show that there exists $R' \in R_1$ such that $(R+R')||_{\gamma_1^X} = \emptyset$.

To construct R' we first notice that $R \in R_0$ and apply proposition 16 —up to a translation $T_{K+L,0}$— to obtain $S \in R_0$ such that

$$S \subset \mathcal{A}_g^0|_{\gamma_{-\infty}^{\infty}}, \quad (S+R)||_{\gamma_{-M}^M} = \emptyset, \quad \gamma_a^b := \square_{K+L+a,0}^{K+L+b,K}. \tag{123}$$

Setting $S' := S|_{\gamma_2^X}$ we get $(S'+R)||_{\gamma_{-M}^M} = \emptyset$ and

$$\mathrm{Sup}(S') \subset \gamma_3^X \cup \gamma_4^X. \tag{124}$$

From proposition 14, corollary 11 and the involved geometry it follows the existence of $S_3, S_4 \subset \mathbf{P}_0(\mathcal{A}_0)$ such that

$$\mathrm{Sup}(S_i) \subset \mathrm{Thk}^N \gamma_i^X, \qquad i = 3,4, \tag{125}$$

and $S' + S_3 + S_4 \in R_0$. We can take $R' := S' + S_1 + S_2$ because $R'|_{\gamma_6^X} = R'$ and $(R+R')||_{\gamma_{-M}^M} = \emptyset$.

2) $\mathcal{A}_1$, R_g^1 and γ_1 satisfy the conditions of proposition 17 and we get $\mathcal{A}_1$ setting $i=1$ in (117). We claim that

$$R_g^2 := T^{(D)}\,\{S \in R_2 \,:\, \mathrm{Sup}(S) \subset \gamma_2\} \tag{126}$$

generates R_2. Indeed, since

$$\Sigma = T^{(D)}\,(\gamma_0 \cup \gamma_1 \cup \gamma_2), \tag{127}$$

for any $S \in R_2$ we have $\mathrm{Sup}(S) \subset T^{(D)}\,(\gamma_2)$. But then

$$R_2 \ni S_{m,n} := S|_{T_{m,n}\,(\gamma_2)}, \tag{128}$$

because for any $b \in \mathcal{B}_g$ there is at most one pair $(m,n) \in \mathbf{Z}^2$ such that $\mathrm{Sup}(b) \cap T_{m,n}\,(\gamma_2) \neq \emptyset$.

3) $\mathcal{A}_3$, R_g^2 and γ_2 satisfy the conditions of proposition 17 and we get $\mathcal{A}_3$ setting $i=2$ in (117). Due to (127) we have $R_4 = \{\emptyset\}$ as desired. $\square$

V. CHARGE

This section deals with charges and string operators. The final goal is to show that the charge groups of pairs of LPGs that are the centralizer of each other —as $\mathcal{G}$ and $\mathcal{S}$ in a TSSG— are isomorphic and finite.

A. Charge groups

We can define

$$\Phi^0(\mathcal{A}) := \Phi^0(\mathcal{A}_g) \tag{129}$$

because the choice of the generating set $\mathcal{A}_g$ is immaterial.

Definition 19 *The charge group of $\mathcal{A}$ is*

$$C_{\mathcal{A}} := \frac{\Phi^0(\mathcal{A})}{\mathrm{Com}_{\mathcal{A}}[\mathcal{P}]}. \tag{130}$$

Its elements are the charges of $\mathcal{A}$.

The quotient group is well defined because $\mathrm{Com}_{\mathcal{A}}[\mathcal{P}] \subset \Phi^0(\mathcal{A})$. When $\phi_1, \phi_2 \in c \in C_{\mathcal{A}}$ we write $\phi_1 \sim \phi_2$.

An isomorphism

$$\mathcal{A} \xrightarrow{F} \mathcal{B}, \tag{131}$$

with $\mathcal{A} \subset \mathcal{P}_A$, $\mathcal{B} \subset \mathcal{P}_B$, induces an isomorphism

$$\begin{aligned} F^* \,:\, \Phi^0(\mathcal{A}) &\longrightarrow \Phi^0(\mathcal{B}) \\ \phi &\longmapsto \phi \circ F^{-1} \end{aligned} \tag{132}$$

satisfying, due to (67),

$$F^*[\mathrm{Com}_{\mathcal{A}}[\mathcal{P}_A]] = \mathrm{Com}_{\mathcal{B}}[\mathcal{P}_B]. \tag{133}$$

Thus, F^* maps charges to charges and gives rise to an isomorphism

$$C_F \,:\, C_{\mathcal{A}} \longrightarrow C_{\mathcal{B}}, \tag{134}$$

making the charge group construction a functor from **LPG** to **Ab** —the category of abelian groups—. This is part of the functor we are looking for.

Coarse graining does not affect the group of charges, in the sense that there exists a natural isomorphism

$$C_{\mathcal{A}} \simeq C_{\mathrm{Crs}_l(\mathcal{A})} \tag{135}$$

for any $l \in \mathbf{N}^*$. It is also clear that the composition of two *LPGs* has a charge group

$$C_{\mathcal{A}\otimes\mathcal{B}} \simeq C_{\mathcal{A}} \times C_{\mathcal{B}}, \tag{136}$$

where the isomorphism is again natural.

B. Charge of generators

Definition 20 *Given an independent set of generators $\mathcal{A}_g$ of a LPG, we say that $a \in \mathcal{A}_g - \langle i\mathbf{1} \rangle$ has charge $c \in C_{\mathcal{A}}$ if*

$$\mathrm{Mor}_{\mathcal{A}_g}(\{a\}) \in c. \tag{137}$$

Notice that the charge of a depends on $\mathcal{A}_g$. Given $\phi \in \Phi^0(\mathcal{A})$, its charge is the product of the charges of the generators in $\overline{\mathrm{Ker}_{\mathcal{A}_g}}(\phi)$.

Proposition 21 *Let $\mathcal{A}$ be a LPG with a LTI set of independent generators $\mathcal{A}_g$ of step L. Let $a_1, a_2 \in \mathcal{A}_g - \langle i\mathbf{1}\rangle$ have respectively charge $c_1, c_2 \in C_{\mathcal{A}}$ and satisfy $a_2 \notin T^{(L)}(\{a_1\})$. Then*

$$\mathcal{A}'_g := \mathcal{A}_g \cup T^{(L)}(\{a_1a_2\}) - T^{(L)}(\{a_1\}) \tag{138}$$

is a LTI set of independent generators of $\mathcal{A}$ for which a_1a_2 has charge c_1 and a_2 has charge c_1c_2.

Proof — Since $a_1a_2 \notin \mathcal{A}_g$ because $\mathcal{A}_g$ is independent, we have $a_2, a_1a_2 \in \mathcal{A}'_g$. Then $a_1 \in \langle \mathcal{A}'_g\rangle$ and by translational invariance $\mathcal{A} = \langle \mathcal{A}'_g\rangle$. Given $S' \in \mathrm{Cnstr}^0(\mathcal{A}'_g)$, let

$$z := \{(i,j) \in \mathbf{Z}^2 \,:\, T_{iL,jL}(a_1a_2) \in S'\}. \tag{139}$$

Then we can construct $S \in \mathrm{Cnstr}^0(\mathcal{A}_g)$ setting

$$S := S' + \sum_{(i,j)\in z} T_{iL,jL}(\{a_1, a_2\}) - \sum_{(i,j)\in z} T_{iL,jL}(\{a_1a_2\}). \tag{140}$$

But $|S| \geq |S'|$ because $T^{(L)}(\{a_1\}) \cap S' = \emptyset$, and thus $\mathcal{A}'_g$ is independent. Let

$$\phi_i = \mathrm{Mor}_{\mathcal{A}_g}(\{a_i\}) \in c_i \in C_{\mathcal{A}}, \qquad i = 1, 2. \tag{141}$$

Then

$$\overline{\mathrm{Ker}_{\mathcal{A}'_g}}(\phi_1) = \{a_1a_2\}, \qquad \overline{\mathrm{Ker}_{\mathcal{A}'_g}}(\phi_2) = \{a_1a_2, a_2\}, \tag{142}$$

and thus

$$\mathrm{Mor}_{\mathcal{A}'_g}(\{a_1a_2\}) = \phi_1, \qquad \mathrm{Mor}_{\mathcal{A}'_g}(\{a_2\}) = \phi_1\phi_2. \tag{143}$$

□

C. Coarse graining

Proposition 22 *Consider a set of LPGs $\bar{\mathcal{A}}_1, \dots, \bar{\mathcal{A}}^n$, on a given qubit lattice, all admiting LTI sets of independent generators and with finite charge groups. There exists $L \in \mathbf{N}^*$ such that the coarse grained LPGs*

$$\mathcal{A}^k = \mathrm{Crs}_L(\bar{\mathcal{A}}^k) \subset \mathcal{P}, \qquad k = 1, \dots, n, \tag{144}$$

for any independent set of generators $\{c_l^k\}_{l=1}^{m_k}$ of $C_{\mathcal{A}^k}$, $m_k \in \mathbf{N}_0$, admit LTI sets of independent generators $\mathcal{A}_g^k$ such that the properties 1 and 2 in proposition 3 and the following ones are satisfied.

3. For any $c \in C_{\mathcal{A}^k}$ there exists $\phi \in c$ such that

$$\mathrm{Sup}_{\mathcal{A}_g^k}(\phi) \subset \{\mathbf{0}\}. \tag{145}$$

4. For any $\phi \in \Phi^0(\mathcal{A}^k)$ and $i, j \in \mathbf{Z}$

$$\phi \sim T_{i,j}(\phi). \tag{146}$$

5. For any $\phi \in \mathrm{Com}_{\mathcal{A}^k}[\mathcal{P}]$ and connected set of sites $\gamma \in \mathbf{P}_0(\Sigma)$ with

$$\mathrm{Sup}_{\mathcal{A}_g^k}(\phi) \subset \gamma, \tag{147}$$

there exists $p \in \mathcal{P}$ such that

$$\mathrm{Com}_{\mathcal{A}^k}(p) = \phi, \qquad \mathrm{Sup}(p) \subset \mathrm{Thk}(\gamma). \tag{148}$$

6. For any $A \in \mathbf{P}_0(\mathcal{A}_g^k)$ and $\gamma = \square_{i,j}^{i+L,j+L}$, where $i, j \in \mathbf{Z}$ and $L \in \mathbf{N}_0$,

$$\mathrm{Sup}(\mathrm{Pro}(A)) \subset \gamma \quad \Longrightarrow \quad \mathrm{Sup}(A) \subset \mathrm{Thk}(\gamma). \tag{149}$$

7. There exist $a_l^k \in \mathcal{A}_g^k$ with charge c_l^k and

$$\mathrm{Sup}(a_l^k) = \{\mathbf{0}\}. \tag{150}$$

Proof — We show, for properties $x = 3, \dots, 7$, that if properties 1 to $x-1$ are satisfied then property x is also satisfied by further coarse graining or changing the sets of generators. Notice that all properties are preserved under coarse graining.

3) Due to the finiteness of $C_{\mathcal{A}^k}$, there exist $m \in \mathbf{N}^*$ such that for any k and $c \in C_{\mathcal{A}^k}$ there exists $\phi \in c$ with

$$\mathrm{Sup}_{\mathcal{A}_g^k}(\phi) \subset \square_{0,0}^{m,m}. \tag{151}$$

Then the LPGs $\mathrm{Crs}_m(\mathcal{A}^k)$ satisfy property 3.

4) The key observation here is that for any $\phi, \phi' \in \Phi^0(\mathcal{A}^k)$, $i, j \in \mathbf{Z}$, we have

$$\phi \sim \phi' \quad \Longleftrightarrow \quad T_{i,j}(\phi) \sim T_{i,j}(\phi'). \tag{152}$$

For each k and $c \in C_{\mathcal{A}^k}$, choose $\phi_c^k \in \Phi^0(\mathcal{A}^k)$ such that

$$\mathrm{Sup}_{\mathcal{A}_g^k}(\phi_c^k) \subset \{\mathbf{0}\}. \tag{153}$$

Due to the finiteness of $C_{\mathcal{A}^k}$, there exist $m_1, m_2 \in \mathbf{N}^*$, $m_1 < m_2$, such that for any k, $c \in C_{\mathcal{A}^k}$ and $i, j \in \mathbf{Z}$

$$T_{m_1,0}(\phi_c^k) \sim T_{m_2,0}(\phi_c^k). \tag{154}$$

Or equivalently, $\phi_c^k \sim T_{m,0}(\phi_c^k)$ with $m = m_2 - m_1$. It follows that for any $\phi \in c \in C_{\mathcal{A}^k}$ we have

$$\phi \sim \phi_c^k \sim T_{m,0}(\phi_c) \sim T_{m,0}(\phi). \tag{155}$$

The same reasoning in the other axis gives another integer $m' \in \mathbf{N}^*$, and the LPGs $\mathrm{Crs}_{mm'}(\mathcal{A}^k) = \langle \mathrm{Crs}_{mm'}(\mathcal{A}_g^k)\rangle$ satisfy properties 1-4.

5) For each k and for each $\phi \in \mathrm{Com}_{\mathcal{A}^k}[\mathcal{P}]$ with $\mathrm{Sup}_{\mathcal{A}_g^k}(\phi) \subset \square_{0,0}^{3,3}$, choose $m' \in \mathbf{N}_0$ such that there exists $p \in \mathcal{P}$ with

$$\mathrm{Com}_{\mathcal{A}^k}(p) = \phi, \qquad \mathrm{Sup}(p) \subset \mathrm{Thk}^{m'}(\square_{0,0}^{3,3}). \tag{156}$$

Let $m-1$ equal the largest of such integers m'. We will show that for any k, $\phi \in \mathrm{Com}_{\mathcal{A}^k}[\mathcal{P}]$ and $\gamma \in \mathbf{P}_0(\Sigma)$ with $\mathrm{Sup}_{\mathcal{A}^k_g}(\phi) \subset \gamma$ and γ connected, there exists $p \in \mathcal{P}$ such that

$$\mathrm{Com}_{\mathcal{A}^k}(p) = \phi, \quad \mathrm{Sup}(p) \subset \mathrm{Thk}^m(\gamma). \tag{157}$$

Then the LPGs $\mathrm{Crs}_m(\mathcal{A}^k) = \langle \mathrm{Crs}_m(\mathcal{A}^k_g)\rangle$ satisfy properties 1-5.

Let $\sigma \in \Sigma$ be such that the set

$$\gamma' := \gamma - \{\sigma\} \tag{158}$$

is connected. There is always such a site if $\gamma \neq \emptyset$. If $|\gamma| = 1$, just notice that $||\mathrm{Thk}(\gamma)|| = 3$ and

$$\mathrm{Thk}^m(\gamma) = \mathrm{Thk}^{m-1}(\mathrm{Thk}(\gamma)). \tag{159}$$

In other case, choose $\sigma' \in \Sigma$ adjacent to σ. Set

$$\phi_0 := \mathrm{Mor}_{\mathcal{A}^k_g}(\overline{\mathrm{Ker}_{\mathcal{A}^k_g}}(\phi)||_{\{\sigma\}}) \tag{160}$$

and notice that

$$\mathrm{Sup}_{\mathcal{A}^k_g}(\phi_0) \subset \gamma_0 := \mathrm{Thk}(\{\sigma\}) \tag{161}$$

due to the second property of proposition 3. Choose $\phi_1 \in \Phi^0(\mathrm{Crs}_{l_2}(\mathcal{A}^k))$ and $p \in \mathcal{P}$ with $\phi_0 \sim \phi_1$ and

$$\begin{gathered} \mathrm{Sup}_{\mathcal{A}^k_g}(\phi_1) \subset \{\sigma'\} \\ \mathrm{Com}_{\mathcal{A}^k}(p) = \phi_0\phi_1, \\ \mathrm{Sup}(p) \subset \mathrm{Thk}^{m-1}(\gamma_0). \end{gathered} \tag{162}$$

This is always possible because $\sigma' \in \mathrm{Thk}(\{\sigma\})$. The result follows by induction on $|\gamma|$ observing that

$$\begin{aligned} &\mathrm{Sup}_{\mathcal{A}^k_g}(\phi\phi_0\phi_1) \subset \gamma', \\ &\mathrm{Thk}^{m-1}(\gamma_0) \cup \mathrm{Thk}^m(\gamma') = \mathrm{Thk}^m(\gamma). \end{aligned} \tag{163}$$

6) We will show that for any k, $A \in \mathbf{P}_0(\mathcal{A}^k_g)$ and $\gamma = \square_{i,j}^{i+L,j+L}$, where $i,j \in \mathbf{Z}$ and $L \in \mathbf{N}_0$,

$$\mathrm{Sup}(\mathrm{Pro}(A)) \subset \gamma \quad \Longrightarrow \quad \mathrm{Sup}(A) \subset \mathrm{Thk}^2(\gamma). \tag{164}$$

Then the LPGs $\mathrm{Crs}_2(\mathcal{A}^k) = \langle \mathrm{Crs}_2(\mathcal{A}^k_g)\rangle$ satisfy properties 1-6.

Let $a \in A$ satisfy

$$\mathrm{Sup}(a) \not\subset \mathrm{Thk}^2(\square_{i,j}^{i+L,j+L}). \tag{165}$$

W.l.o.g, we assume that $\mathrm{Sup}(a) \subset \square_{u,v}^{u+2,v+2}$ with $u \geq i+1+L$. Choose $w \in \mathbf{N}^*$ such that $T_{w,0}(a) \notin A$ and $p \in \mathcal{P}$ such that $\mathrm{Sup}(p) \subset \mathrm{Thk}(\square_{u,v}^{u+w+2,v+2})$ and

$$\mathrm{Com}_{\mathcal{A}^k}(p) = \phi\, T_{w,0}(\phi), \quad \phi := \mathrm{Mor}_{\mathcal{A}^k_g}(\{a\}). \tag{166}$$

Then $\mathrm{Sup}(p) \cap \mathrm{Sup}(\mathrm{Pro}(A)) = \emptyset$ implies $(p;A) = 1$, but at the same time $(p;A) = \phi(A) = -1$, a contradiction.

7) For each k, let us show that there exist a list of LTI sets of independent generators

$$\mathcal{A}^k_g = \mathcal{A}^{k,0}_g, \mathcal{A}^{k,1}_g, \ldots, \mathcal{A}^{k,m_k}_g = \tilde{\mathcal{A}}^k_g \tag{167}$$

such that for any $t = 1,\ldots,m_k$ and any $l = 1,\ldots,t$ there exists $a_l \in \mathcal{A}^{k,t}_g$ with charge c^k_l and satisfying (150). Moreover, we do it in such a way that all these generating sets preserve properties 1-6, so that the final generating sets $\tilde{\mathcal{A}}^k_g$ satisfy all the desired properties.

We construct $\mathcal{A}^{k,t}_g$ from $\mathcal{A}^{k,t-1}_g$ as follows. There exists $\phi \in c^k_t$ with $\mathrm{Sup}_{\mathcal{A}^{k,t-1}_g}(\phi) = \{\mathbf{0}\}$. Also, there exists $\hat{a}_0 \in \overline{\mathrm{Ker}_{\mathcal{A}^{k,t-1}_g}}(\phi)$ such that $\hat{a}_0 \neq a_k$ for any $1 \leq k < t$, because the charge c^k_t is independent of the charges $c^k_1, \ldots, c^k_{t-1}$. Label the elements of $\overline{\mathrm{Ker}_{\mathcal{S}^{k,t-1}_g}}(\phi) - \{\hat{a}_0\}$ as $\{\hat{a}_i\}_{i=1}^r$ and apply proposition 21 repeatedly to perform the substitutions $\hat{s}_i \to \hat{a}_0\hat{a}_i$, $i = 1,\ldots,r$. Let $\mathcal{A}^{k,t}_g$ be the resulting set of generators. Then $\hat{s}_0$ has charge c^k_t in $\mathcal{A}^{k,t}_g$ and the rest of generators with support in $\mathbf{0}$ have in $\mathcal{A}^{k,t}_g$ the same charge as in $\mathcal{A}^{k,t-1}_g$. □

D. Strings

Let $\mathcal{A}$ satisfy the properties in proposition 22. We define the following sets of 'string operators'.

Definition 23 *Given $\phi \in \Phi^0(\mathcal{A})$, $\gamma \subset \Sigma$, we set*

$$\begin{aligned} \mathrm{Str}(\phi;\gamma) := \{p \in \mathcal{P} \,:\, &\mathrm{Com}_{\mathcal{A}_g}(p) = \phi \,\wedge \\ &\wedge\ \mathrm{Sup}(p) \subset \mathrm{Thk}(\gamma)\}. \end{aligned} \tag{168}$$

Given $\phi_1, \phi_2 \in \Phi^0(\mathcal{A})$ with

$$\mathrm{Sup}_{\mathcal{A}_g}(\phi_i) = \{\sigma_i\} \subset \Sigma, \qquad i = 1,2, \tag{169}$$

we set

$$\begin{aligned} \mathrm{Str}(\phi_1,\phi_2) &:= \mathrm{Str}(\phi_1\phi_2; \mathrm{Path}(\sigma_1,\sigma_2)), \\ \mathrm{Str}(1,\phi_1) &:= \mathrm{Str}(\phi_1;1) := \mathrm{Str}(\phi_1;\{\sigma_1\}), \\ \mathrm{Str}(1;1) &:= \{\mathbf{1}\}. \end{aligned} \tag{170}$$

Finally, we set for $c \in C_{\mathcal{A}}$ and a path $\gamma := (\sigma_i)_{i=1}^n$,

$$\begin{aligned} \mathrm{Str}(c;\gamma) := \bigcup \{&\mathrm{Str}(\phi_1\phi_2;\gamma) \,:\, \phi_1,\phi_2 \in c \,\wedge \\ &\wedge\ \mathrm{Sup}(\phi_1) \subset \sigma_1 \ \wedge\ \mathrm{Sup}(\phi_2) \subset \sigma_n\}. \end{aligned} \tag{171}$$

In general $\mathrm{Str}(\phi;\gamma)$ can be empty. However, suppose that for some $n \in \mathbf{N}_0$ we have

$$\begin{aligned} \phi &= \prod_{k=1}^n \phi_k, \qquad \phi_k \in \mathrm{Com}_{\mathcal{A}}[\mathcal{P}], \\ \gamma &= \bigcup_{k=1}^n \gamma_k, \qquad \gamma_k \in \mathbf{P}_0(\Sigma), \end{aligned} \tag{172}$$



FIG. 12: The geometry of proposition 24 for $l = 8$. The striped sites are the common endpoints of two string operators p_1 and p_2, the first running trough the bottom and right side, the second through the top and right side. The central shaded set of sites is $\square_{3,3}^{l-2,l-2}$.

with the sets γ_k connected and

$$\mathrm{Sup}_{\mathcal{A}_g}(\phi_k) \subset \gamma_k, \qquad k = 1, \dots, n. \tag{173}$$

Then due to property 5 $\mathrm{Str}(\phi;\gamma)$ is nonempty. In particular, $\mathrm{Str}(\phi_1, \phi_2)$ is nonempty iff $\phi_1 \sim \phi_2$, and $\mathrm{Str}(c;\gamma)$ is always nonempty.

Proposition 24 *Let $\mathcal{B} = \mathscr{Z}_{\mathcal{P}}(\mathcal{A})$, $l \in \mathbf{N}^*$. Let $\phi \in c \in C_{\mathcal{A}}$ for c nontrivial and $p_1, p_2 \in \mathcal{P}$, $B \in \mathbf{P}_0(\mathcal{B}_g)$ be such that, see Fig. 12,*

$$\begin{aligned} \mathrm{Sup}_{\mathcal{A}_g}(\phi) &= \{\mathbf{0}\}, \\ p_1 &\in \mathrm{Str}(\phi, T_{l,l}(\phi)), \\ p_2 &\in \mathrm{Str}(T_{l,l}(\phi), \phi), \\ p_1 p_2 &\in \mathrm{Pro}(B). \end{aligned} \tag{174}$$

Then

$$\square_{3,3}^{l-2,l-2} \subset \mathrm{Sup}(B). \tag{175}$$

Proof — It is enough to consider $l \geq 6$. Let $\gamma := \square_{3,3}^{l-2,l-2}$ and suppose that there is a site $(i,j) \in \gamma - \mathrm{Sup}(B)$. Define the sets of sites

$$\begin{aligned} \gamma_1 &:= \square_{i,j}^{i+1,\infty}, \\ \gamma_2 &:= \square_{-\infty,-\infty}^{i,\infty}, \\ \gamma_3 &:= \square_{i-1,-1}^{i+2,3}. \end{aligned} \tag{176}$$

Choose $b \in \mathrm{Pro}(B||_{\gamma_1})$ and set $p_3 := b p_1 p_2$, so that

$$\mathrm{Sup}(b) \subset \square_{i-1,j}^{i+2,\infty}, \qquad \gamma_1 \cap \mathrm{Sup}(p_3) = \emptyset. \tag{177}$$

Choose $p, q, r \in \mathcal{P}$ with

$$p \propto p_1|_{\gamma_2}, \qquad q \propto p_3|_{\gamma_2}. \tag{178}$$

Let $\Sigma_2 := \{\square_{a,b}^{a+2,b+2} : (a,b) \in \Sigma\}$. Since $p_3 \in \mathscr{Z}_{\mathcal{P}}(\mathcal{A})$,

$$\begin{aligned} \mathrm{Sup}_{\mathcal{A}_g}(q) \subset \bigcup \{\gamma \in \Sigma_2 \,:\, &\mathrm{Sup}(p_3) \cap \gamma_2 \cap \gamma \neq \emptyset \,\wedge \\ &\wedge\, \mathrm{Sup}(p_3) \cap (\Sigma - \gamma_2) \cap \gamma \neq \emptyset\} \subset \gamma_3. \end{aligned} \tag{179}$$

Let $\phi_1 \in c$ be such that

$$\mathrm{Com}_{\mathcal{A}}(p) = \phi_1\, \phi. \tag{180}$$

Then

$$\begin{aligned} \mathrm{Sup}_{\mathcal{A}_g}(\phi_1) \subset \bigcup \{\gamma \in \Sigma_2 \,:\, &\mathrm{Sup}(p) \cap \gamma_2 \cap \gamma \neq \emptyset \,\wedge \\ &\wedge\, \mathrm{Sup}(p) \cap (\Sigma - \gamma_2) \cap \gamma \neq \emptyset\} \cup \\ &\cup (\mathrm{Sup}(T_{l,l}(\phi)) \cap \gamma_2) \subset \gamma_3. \end{aligned} \tag{181}$$

But from

$$\mathrm{Thk}(\gamma_3) \cap (\mathrm{Sup}(b) \cup \mathrm{Sup}(p_2)) = \emptyset \tag{182}$$

we get $\mathrm{Sup}_{\mathcal{A}_g}(q) = \mathrm{Sup}_{\mathcal{A}_g}(q)|_{\gamma_3} = \mathrm{Sup}_{\mathcal{A}_g}(p)|_{\gamma_3}$. Thus $\mathrm{Com}_{\mathcal{A}_g}(q) = \phi \in c$, a contradiction. $\square$

E. Charge and constraints

Proposition 25 *Let $\mathcal{A}_g$ be a LTI set of independent generators of $\mathcal{A}$ with $\mathrm{Cnstr}(\mathcal{A}_g)$ finite. $C_{\mathcal{A}}$ admits an independent set of generators C_g such that the sets*

$$R_c := \bigcup_{c' \in \bar{c}} A_{c'}, \qquad c \in C_g, \tag{183}$$

form a set of independent generators of $\mathrm{Cnstr}(\mathcal{A}_g)$, where for $c \in C_g$

$$\bar{c} := C_{\mathcal{A}} - \langle C_g - \{c\} \rangle \tag{184}$$

$$A_c := \{a \in \mathcal{A}_g - \langle i\mathbf{1} \rangle \,:\, \mathrm{Mor}_{\mathcal{A}_g}(\{a\}) \in c\}. \tag{185}$$

Proof — Consider the set of nontrivial charges such that there exist generators with that charge

$$C := \{c \in C_{\mathcal{A}} - 1 \,:\, \exists a \in \mathcal{A}_g - \langle i\mathbf{1} \rangle \quad \mathrm{Mor}_{\mathcal{A}_g}(\{a\}) \in c\}. \tag{186}$$

Clearly $C_{\mathcal{A}} = \langle C \rangle$ and we can choose C_g to be a possibly infinite, but countable, subset of C.

For any $c \in C$ and $R \in \mathrm{Cnstr}(\mathcal{A}_g)$, either $A_c \cap R = \emptyset$ or $A_c \cap R = A_c$. In other case, there exist $a_1, a_2 \in A_c$ with $a_1 \in R$, $a_2 \notin R$. But $\phi_1 \sim \phi_2$ for $\phi_i := \mathrm{Mor}_{\mathcal{A}_g}(\{a_i\})$. Thus, there exists $p \in \mathcal{P}$ with $\mathrm{Com}_{\mathcal{A}}(p) = \phi_1 \phi_2$ and thus $(p; R) = -1$, a contradiction.

Given $c_0 \in C_g$, let us show that $R_{c_0} \in \mathrm{Cnstr}(\mathcal{A}_g)$. Take any $p \in \mathcal{P}$ and let

$$C_p := \{c \in C \,:\, (p; A_c) = -1\}. \tag{187}$$

Set $A_p := \mathrm{Sup}_{\mathcal{A}_g}(p)$ and for any $a \in \mathcal{A}_g$ let c_a be the charge a. Since i) $\mathrm{Com}_{\mathcal{A}}(p)$ has trivial charge, ii) $(p; A_c) = -1$ if and only if $|A_c \cap A_p|$ is odd, and iii) $c^2 = 1$ for any $c \in C$, we get

$$1 = \prod_{a\in A_p} c_a = \prod_{c\in C} c^{|A_c\cap A_p|} = \prod_{c\in C_p} c. \qquad (188)$$

For each $c \in \overline{c_0}$ write $c = c_1 c_0$, so that $c_1 \in \langle C_g - \{c_0\}\rangle$. Then

$$1 = \prod_{c\in C_p} c = c' {c_0}^{|C_p\cap\overline{c_0}|}, \qquad c' \in \langle C_g - \{c_0\}\rangle. \qquad (189)$$

If $|C_p \cap \overline{c_0}|$ were odd this would give $c_0 \in \langle C_g - \{c_0\}\rangle$, a contradiction. Therefore

$$(p; R_{c_0}) = (-1)^{|C_p\cap\overline{c_0}|} = 1, \qquad (190)$$

showing that indeed $R_{c_0} \in \mathrm{Cnstr}(\mathcal{A}_g)$.

The global constraints $\{R_c\}$ are independent since for any $c, c' \in C_g$ we have $A_c \subset R_{c'} \iff c = c'$. Due to lemma 15, C_g is finite. Thus it is enough to show that for any $R \in \mathrm{Cnstr}(\mathcal{A}_g)$

$$R \cap \bigcup_{c\in C_g} A_c = \emptyset \Rightarrow R = \emptyset. \qquad (191)$$

So let R satisfy the LHS and suppose that there exists $c_0 \in C - C_g$ such that $A_{c_0} \subset R$. There exists a unique $C_R \subset C_g$ such that $c_0 = \prod_{c\in C_R} c$. For each $c \in C_g$, choose $a_c \in A_c$. Then there exists $p \in \mathcal{P}$ with

$$\mathrm{Com}_{\mathcal{A}}(p) = \mathrm{Mor}_{\mathcal{A}_g}(\{a_{c_0}\} \cup \bigcup_{c\in C_R} \{a_c\}). \qquad (192)$$

But $(p; R) = (p; a_{c_0}) = -1$, a contradiction. ☐

Corolary 26 *There exists $n \in \mathbf{N}_0$ such that*

$$C_{\mathcal{A}} \simeq \mathrm{Cnstr}(\mathcal{A}_g) \simeq \mathbf{Z}_2^n. \qquad (193)$$

Proposition 27 *Let $\mathcal{A}, \mathcal{B}$ be LPGs with $C_{\mathcal{A}}$ finite and $\mathcal{B} = \langle \mathcal{B}_g \rangle = \mathcal{Z}_{\mathcal{P}}(\mathcal{A})$. $C_{\mathcal{A}}$ is isomorphic to a subgroup of* $\mathrm{Cnstr}(\mathcal{B}_g)$.

Proof — W.l.o.g. let $\mathcal{A}_g$ satisfy the properties in proposition 22 and $\mathcal{B}_g$ have step 1. Let $\{c_k\}_{k=1}^n$ be a set of independent generators of $C_{\mathcal{A}}$. For each $k = 1, \dots, n$, choose $\phi_k \in c_k \in C_{\mathcal{A}}$ and $h_k, v_k \in \mathcal{P}$, $B_k \in \mathbf{P}_0(\mathcal{B}_g)$ such that

$$\begin{aligned} \mathrm{Sup}_{\mathcal{A}_g}(\phi_k) &= \mathbf{0}, \\ h_k &\in \mathrm{Str}(\phi_k, T_{1,0}\,(\phi_k)), \\ v_k &\in \mathrm{Str}(\phi_k, T_{0,1}\,(\phi_k)), \\ \mathrm{Pro}(B_k) &\ni h_k\, T_{0,1}\,(h_k)\, v_k\, T_{1,0}\,(v_k), \end{aligned} \qquad (194)$$

For $l \in \mathbf{N}^*$, let

$$\begin{aligned} h_k^l &:= T_{0,0}\,(h_k) T_{1,0}\,(h_k) \cdots T_{l-1,0}\,(h_k), \\ v_k^l &:= T_{0,0}\,(v_k) T_{0,1}\,(v_k) \cdots T_{0,l-1}\,(v_k), \end{aligned} \qquad (195)$$

Any $c \in C_{\mathcal{A}}$ can be expressed as

$$c = \prod_{j=1}^n c_j^{x_j}, \quad x_j = 0, 1, \quad j = 1, \dots, n. \qquad (196)$$

Set $\phi_c := \phi_1^{x_1} \cdots \phi_n^{x_n} \in c$ and

$$\begin{aligned} h_c^l &:= (h_1^l)^{x_1} \cdots (h_n^l)^{x_n}, \\ v_c^l &:= (v_1^l)^{x_1} \cdots (v_n^l)^{x_n}, \\ B_c^l &:= \sum_{i,j=0}^{l-1} \sum_{k=1}^n T_{i,j}\,(B_k^{x_k}), \end{aligned} \qquad (197)$$

so that

$$\begin{aligned} h_c^l &\in \mathrm{Str}(\phi_c, T_{l,0}\,(\phi_c)), \\ v_c^l &\in \mathrm{Str}(\phi_c, T_{0,l}\,(\phi_c)), \\ \mathrm{Pro}(B_c^l) &\ni h_c^l\, T_{0,l}\,(h_c^l)\, v_c^l\, T_{l,0}\,(v_c^l), \end{aligned} \qquad (198)$$

Consider the morphism $f : C_{\mathcal{A}} \longrightarrow \mathrm{Cnstr}(\mathcal{B}_g)$ defined for $c \in C_{\mathcal{A}}$ by $f(c) = R_c$ with

$$R_c := T^{(1)}\,(B_c^1) \in \mathrm{Cnstr}(\mathcal{B}_g). \qquad (199)$$

Let $R_c = \emptyset$ for some $c \in C_{\mathcal{A}}$. Choose $l \geq 2$ such that $\mathrm{Sup}(B_c^1) \subset \mathrm{Thk}^l(\square_{0,0}^{2,2})$. Then for $L := 2l + 2$ we have

$$\mathrm{Sup}(R_c + B_c^L) \subset \bigcup_{i,j\in I} \mathrm{Sup}(T_{i,j}\,(B_c^1)) \not\ni (l+1, l+1) \qquad (200)$$

with $I = \mathbf{Z} - \{0, \dots, L\}$. Thus $B_c^L|_{\{(l+1,l+1)\}} = \emptyset$, which implies that c is trivial using proposition 24, showing that f is a monomorphism. ☐

Corolary 28 *Let $\mathcal{A}, \mathcal{B}$ be LPGs with $\mathcal{A} = \mathcal{Z}_{\mathcal{P}}(\mathcal{B})$ and $\mathcal{B} = \mathcal{Z}_{\mathcal{P}}(\mathcal{A})$. Then*

$$C_{\mathcal{A}} \simeq C_{\mathcal{B}}. \qquad (201)$$

Proof — This follows from corollary 26 and proposition 27, choosing $\mathcal{A}_g$ and $\mathcal{B}_g$ independent, which is possible according to theorem 18. ☐

VI. TOPOLOGICAL CHARGE

The commutation properties of string operators sometimes only depend on the charges and topology of the involved strings. As shown in this section, this gives rise to invariants that we put together under the label of topological charge.

A. String commutation rules

Proposition 29 *Let $\mathcal{A}, \mathcal{B}$ be LPGs on a given lattice satisfying the properties in proposition 22 with*

$$\mathcal{Z}_{\mathcal{P}}(\mathcal{B}) \subset \mathcal{Z}_{\mathcal{P}}(\mathcal{Z}_{\mathcal{P}}(\mathcal{A})). \qquad (202)$$



FIG. 13: The geometry of proposition 29 for $a_1 = -5$, $a_2 = 4$, $b_1 = -3$ and $b_2 = 4$. The striped sites are the endpoints of two string operators p_A and p_B.

Let $p_A, p_B \in \mathcal{P}$ *be such that, see Fig. 13,*

$$\begin{aligned} p_A &\in \mathrm{Str}(c_A; \mathrm{Path}((a_1, 0), (a_2, 0))), \\ p_B &\in \mathrm{Str}(c_B; \mathrm{Path}((0, b_1), (0, b_2))), \end{aligned} \tag{203}$$

for some integers $a_1, b_1 \leq -3$, $a_2, b_2 \geq 3$, *and* $c_A \in C_{\mathcal{A}}$, $c_B \in C_{\mathcal{B}}$. *The quantity*

$$\kappa(c_A, c_B) := (p_A; p_B), \tag{204}$$

only depends on c_A *and* c_B. *Moreover, for any* $c'_A \in C_{\mathcal{A}}$,

$$\begin{aligned} \kappa(c_A, c_B) &= \kappa(c_B, c_A), \\ \kappa(c_A c'_A, c_B) &= \kappa(c_A, c_B)\kappa(c'_A, c_B). \end{aligned} \tag{205}$$

Proof — Let $\bar{p}_A, \bar{p}_B, \bar{a}_1, \bar{a}_2, \bar{b}_1, \bar{b}_2$ be an alternative choice for the definition of $\kappa(c_A, c_B)$, with the same properties as $p_A, p_B, a_1, a_2, b_1, b_2$. Set $a := a_2 - \bar{a}_1$, $b := b_2 + \bar{b}_2 + 3$, and choose $q_A, q_B \in \mathcal{P}$ such that

$$\begin{aligned} q_A &\in \mathrm{Str}(\mathrm{Com}_{\mathcal{A}}(p_A \bar{p}_A); \gamma_A), \\ q_B &\in \mathrm{Str}(\mathrm{Com}_{\mathcal{B}}(p_B \bar{p}_B); \gamma_B), \end{aligned} \tag{206}$$

where,

$$\begin{aligned} \gamma_A &:= \mathrm{Path}((a_1, 0), (a_1, b), (\bar{a}_1 + a, 0)) \cup \square_{a_2,0}^{a_2+1,1}, \\ \gamma_B &:= \mathrm{Path}((0, b_1), (a, \bar{b}_1)) \cup \mathrm{Path}((0, b_2), (a, \bar{b}_2)). \end{aligned} \tag{207}$$

Then, setting $\tilde{p}_A := T_{a,0}\,(\bar{p}_A)$, $\tilde{p}_B := T_{a,0}\,(\bar{p}_B)$,

$$\begin{aligned} p_A \tilde{p}_A q_A &\in \mathcal{Z}_{\mathcal{P}}(\mathcal{A}), \\ p_B \tilde{p}_B q_B &\in \mathcal{Z}_{\mathcal{P}}(\mathcal{B}), \end{aligned} \tag{208}$$

and for $k, k' = 1, 2$

$$\begin{aligned} \mathrm{Sup}(q_A) \cap (\mathrm{Sup}(q_B) \cup \mathrm{Sup}(p_B) \cup \mathrm{Sup}(\tilde{p}_B)) &= \emptyset, \\ \mathrm{Sup}(q_B) \cap (\mathrm{Sup}(q_A) \cup \mathrm{Sup}(p_A) \cup \mathrm{Sup}(\tilde{p}_A)) &= \emptyset, \\ \mathrm{Sup}(p_A) \cap (\mathrm{Sup}(\tilde{p}_B) &= \emptyset, \\ \mathrm{Sup}(p_B) \cap (\mathrm{Sup}(\tilde{p}_A) &= \emptyset, \end{aligned} \tag{209}$$

which gives

$$\begin{aligned} (p_A; p_B)(\bar{p}_A; \bar{p}_B) &= (p_A; p_B)(\tilde{p}_A; \tilde{p}_B) = \\ &= (p_A \tilde{p}_A q_A^1 q_A^2; p_B \tilde{p}_B q_B^1 q_B^2) = 1. \end{aligned} \tag{210}$$

Regarding the first property in (205), first notice that condition (202) is symmetric, so that $\kappa(c_B, c_A)$ is defined. Choose $\phi_A \in C_{\mathcal{A}}$, $\phi_B \in C_{\mathcal{B}}$ with

$$\mathrm{Sup}(\phi_A), \mathrm{Sup}(\phi_B) \subset \square_{0,0}^{1,1}. \tag{211}$$

Let $p_A, q_A, r_A, p_B, q_B, r_B \in \mathcal{P}$ be such that

$$\begin{aligned} p_A &\in \mathrm{Str}(T_{-6,0}\,(\phi_A), T_{0,0}\,(\phi_A)), \\ q_A &\in \mathrm{Str}(T_{0,-6}\,(\phi_A), T_{0,0}\,(\phi_A)), \\ r_A &\in \mathrm{Str}(T_{0,-6}\,(\phi_A), T_{-6,0}\,(\phi_A)), \\ p_B &\in \mathrm{Str}(T_{3,-3}\,(\phi_B), T_{-3,-3}\,(\phi_B)), \\ q_B &\in \mathrm{Str}(T_{-3,3}\,(\phi_B), T_{-3,-3}\,(\phi_B)), \\ r_B &\in \mathrm{Str}(T_{-3,3}\,(\phi_B), T_{3,-3}\,(\phi_B)), \end{aligned} \tag{212}$$

so that, reasoning as in (210),

$$\begin{aligned} \kappa(c_A, c_B)\kappa(c_B, c_A) &= (p_A; p_B)(q_A; q_B) = \\ &= \kappa(p_A q_A r_A, p_B q_B r_B) = 1. \end{aligned} \tag{213}$$

As for the second property in (205), just notice that

$$\begin{aligned} p_1 \in \mathrm{Str}(c_1; \mathrm{Path}(\sigma, \sigma')) \wedge p_2 \in \mathrm{Str}(c_2; \mathrm{Path}(\sigma, \sigma')) \Longrightarrow \\ \Longrightarrow p_1 p_2 \in \mathrm{Str}(c_1 c_2; \mathrm{Path}(\sigma, \sigma')) \end{aligned} \tag{214}$$

for any $c_1, c_2 \in C_{\mathcal{A}}$ and $\sigma, \sigma' \in \Sigma$. □

Proposition 30 *Let* $\mathcal{A}, \mathcal{B}$ *be as in proposition 29. Let* $c_A \in C_{\mathcal{A}}$ *be a non-trivial charge. There exists* $c_B \in C_{\mathcal{B}}$ *with*

$$\kappa(c_A, c_B) = -1. \tag{215}$$

Proof — Choose $\mathcal{B}_g$ be independent, and $\phi \in c$, $h, v \in \mathcal{P}$, $B \in \mathbf{P}_0(\mathcal{B}_g - \langle i\mathbf{1} \rangle)$ such that

$$\begin{aligned} \mathrm{Sup}_{\mathcal{A}_g}(\phi) &= \mathbf{0}, \\ h &\in \mathrm{Str}(\phi, T_{6,0}\,(\phi)), \\ v &\in \mathrm{Str}(\phi, T_{0,6}\,(\phi)), \\ \mathrm{Pro}(B) &\ni h\, T_{0,6}\,(h)\, v\, T_{6,0}\,(v). \end{aligned} \tag{216}$$

According to proposition 24 we can choose $b \in B$ with $\mathrm{Sup}(b) \subset \square_{3,3}^{4,4}$. Let c_B be the charge of b in $\mathcal{B}_g$ and choose $L \in \mathbf{N}^*$ such that $T_{-L,0}\,(b) \notin B$ and $p \in \mathrm{Str}(\phi; T_{0,-L}\,(\phi))$, $\phi := \mathrm{Mor}_{\mathcal{B}_g}(b)$. Then $\kappa(c_A, c_B) = (p; v) = (p; r) = -1$ for any $r \in \mathrm{Pro}(B)$ because $\mathrm{Sup}(p) \cap \mathrm{Sup}(h T_{0,6}\,(h) T_{6,0}\,(v)) = \emptyset$. □

Proposition 31 *Let* $\mathcal{A}$ *be a LPG satisfying the properties in proposition 22 with*

$$\mathcal{Z}_{\mathcal{P}}(\mathcal{A}) \subset \mathcal{Z}_{\mathcal{P}}(\mathcal{Z}_{\mathcal{P}}(\mathcal{A})). \tag{217}$$



FIG. 14: The geometry of proposition 31 for $a_1 = -4$, $a_2 = 5$, $b_1 = -5$ and $b_2 = 3$. The striped sites are the endpoints of four string operators p_i, each with an endpoint in the central striped site.

Let $c \in C_{\mathcal{A}}$*,* $P = \{p_1, p_2, p_3, p_4\} \subset \mathcal{P}$ *be such that for* $k, l = 1, 2, 3, 4$

$$\begin{aligned} p_k &\in \mathrm{Str}(c; \gamma_k), \\ p_k p_l &\in \mathrm{Str}(c; \gamma_l \circ \gamma_k^{-1}), \end{aligned} \tag{218}$$

where

$$\begin{aligned} \gamma_1 &= \mathrm{Path}(\mathbf{0}, (a_1, 0)), \\ \gamma_2 &= \mathrm{Path}(\mathbf{0}, (a_2, 0)), \\ \gamma_3 &= \mathrm{Path}(\mathbf{0}, (0, b_1)), \\ \gamma_4 &= \mathrm{Path}(\mathbf{0}, (0, b_2)), \end{aligned} \tag{219}$$

for some integers $a_1, b_1 \leq -3$*,* $a_2, b_2 \geq 3$*. Let* $p, q, r \in P$ *be all different. The quantity*

$$\theta(c) := (p; q)(p; r)(q; r) = (pq; pr), \tag{220}$$

only depends on c*. Moreover, for any* $c_1, c_2 \in C_{\mathcal{A}}$*,*

$$\theta(c_1 c_2) = \theta(c_1)\theta(c_2)\kappa(c_1, c_2). \tag{221}$$

Proof — Notice that there exist $\phi_k \in \Phi^0(\mathcal{A})$ such that

$$\begin{aligned} \mathrm{Sup}(\phi_0) &\subset \square_{0,0}^{1,1}, \\ \mathrm{Sup}(\phi_1) &\subset \square_{a_1,0}^{a_1+1,1}, \\ \mathrm{Sup}(\phi_2) &\subset \square_{a_2,0}^{a_2+1,1}, \\ \mathrm{Sup}(\phi_3) &\subset \square_{0,b_1}^{1,b_1+1}, \\ \mathrm{Sup}(\phi_4) &\subset \square_{0,b_2}^{1,b_2+1}, \end{aligned} \tag{222}$$

and

$$p_k \in \mathrm{Str}(\phi_0, \phi_k), \qquad k = 1, 2, 3, 4, \tag{223}$$

Let $\bar\phi_k, \bar p_{k'}, \bar a_1, \bar a_2, \bar b_1, \bar b_2$ be an alternative choice for the definition of $\kappa(c_A, c_B)$, with the same properties as $\bar\phi_k, p_{k'}, a_1, a_2, b_1, b_2$, $k = 0, \ldots, 4$, $k' = 1, \ldots, 4$. We show, for example, that $(p_1 p_4; p_2 p_4) = (\bar p_1 \bar p_3; \bar p_3 \bar p_4)$, the other combinations are similar. Set $b := b_2 - \bar b_1$ and choose $q_1, q_2, q_3 \in \mathcal{P}$ such that

$$\begin{aligned} q_1 &\in \mathrm{Str}(\phi_1 T_{0,b}(\bar\phi_1); \mathrm{Path}((a_1, 0), (\bar a_1, b))), \\ q_2 &\in \mathrm{Str}(\phi_2 T_{0,b}(\bar\phi_4); \mathrm{Path}((0, b + \bar a_2), (a_2, 0))), \\ q_3 &\in \mathrm{Str}(\phi_4 T_{0,b}(\bar\phi_3); \mathrm{Path}((0, b_2), (0, b + \bar b_1))). \end{aligned} \tag{224}$$

Then, reasoning as in (210)

$$\begin{aligned} &(p_1 p_4; p_2 p_4)(T_{0,b}(\bar p_1 \bar p_3); T_{0,b}(\bar p_3 \bar p_4)) = \\ &= (p_1 p_4 T_{0,b}(\bar p_1 \bar p_3) q_1 q_3; p_2 p_4 T_{0,b}(\bar p_3 \bar p_4) q_2 q_3) = 1. \end{aligned} \tag{225}$$

We next prove (221). Choose $\phi_1 \in c_1, \phi_2 \in c_2$ with

$$\mathrm{Sup}(\phi) \cup \mathrm{Sup}(\phi') \subset \square_{0,0}^{1,1}. \tag{226}$$

Let $p, q, r, p', q', r' \in \mathcal{P}$ be such that

$$\begin{aligned} p &\in \mathrm{Str}(T_{0,0}(\phi_1), T_{0,-11}(\phi_1)), \\ q &\in \mathrm{Str}(T_{0,0}(\phi_1), T_{12,0}(\phi_1)), \\ r &\in \mathrm{Str}(T_{0,0}(\phi_1), T_{0,12}(\phi_1)), \\ p' &\in \mathrm{Str}(T_{3,3}(\phi_2), T_{3,-8}(\phi_2)), \\ q' &\in \mathrm{Str}(T_{3,3}(\phi_2), T_{15,3}(\phi_2)), \\ r' &\in \mathrm{Str}(T_{3,3}(\phi_2), T_{3,15}(\phi_2)). \end{aligned} \tag{227}$$

Then

$$\begin{aligned} &(pp'qq'; pp'rr') = (pq; qr)(p'q'; q'r')(p'; q) = \\ &\qquad = \theta(c)\theta(c')\kappa(c,)c'. \end{aligned} \tag{228}$$

But, setting $\phi := \mathrm{Crs}_4(\phi_1 T_{3,3}(\phi_2))$,

$$\begin{aligned} \mathrm{Crs}_4(pp') &\in \mathrm{Str}(T_{0,0}(\phi), T_{0,-3}(\phi)), \\ \mathrm{Crs}_4(qq') &\in \mathrm{Str}(T_{0,0}(\phi), T_{3,0}(\phi)), \\ \mathrm{Crs}_4(qq') &\in \mathrm{Str}(T_{0,0}(\phi), T_{0,3}(\phi)), \end{aligned} \tag{229}$$

and, using the observation (233) below, we get

$$\begin{aligned} &\kappa(c, c') = \kappa(\mathrm{Crs}_4(c), \mathrm{Crs}_4(c)) = \\ &\quad = (\mathrm{Crs}_4(pp')\mathrm{Crs}_4(qq'); \mathrm{Crs}_4(pp')\mathrm{Crs}_4(rr')) = \\ &\quad (pp'qq'; pp'rr') = (pq; qr)(p'q'; q'r')(p'; q) = \\ &\qquad = \theta(c)\theta(c')\kappa(c, c'). \end{aligned} \tag{230}$$

□

Clearly

$$\kappa(c, 1) = \theta(1) = 1 \tag{231}$$

with 1 denoting the trivial charge. Then (221) implies

$$\kappa(c, c) = 1. \tag{232}$$

B. Invariance

Notice that κ and θ are invariant under coarse graining, up to the natural isomorphism (135),

$$\kappa = \kappa \circ (\mathrm{Crs}_l \times \mathrm{Crs}_l), \qquad \theta = \theta \circ \mathrm{Crs}_l, \tag{233}$$

because for any $c \in C_{\mathcal{A}}$, path γ and $l \in \mathbf{N}^*$

$$\mathrm{Crs}_l(\mathrm{Str}(c;\gamma)) \subset \mathrm{Str}(\mathrm{Crs}_l(c); \mathrm{Crs}_l(\gamma)). \tag{234}$$

As for the behavior under LPG composition, consider the projections

$$p_i : C_{\mathcal{A}_1 \otimes \mathcal{A}_2} \longrightarrow C_{\mathcal{A}_i}, \qquad i = 1,2, \tag{235}$$

constructed according to the isomorphism (136). Clearly, when both sides make sense,

$$\begin{aligned} \kappa &= (\kappa \circ (p_1 \times p_1))(\kappa \circ (p_2 \times p_2)), \\ \theta &= (\kappa \circ p_1)(\kappa \circ p_2). \end{aligned} \tag{236}$$

Finally, given an LPG isomorphism F from $\mathcal{A}$ to $\mathcal{B}$ that can be regarded also as an LPG isomorphism F' from $\mathcal{A}'$ to $\mathcal{B}'$, whenever both sides make sense we have,

$$\kappa = \kappa \circ (C_F \times C_{F'}), \qquad \theta = \theta \circ C_F, \tag{237}$$

because, for example, for any $c \in C_{\mathcal{A}}$, $L \in \mathbf{Z}$,

$$\mathrm{Crs}_{2r+1}(F[\mathrm{Str}(c;\gamma_1)]) \subset \mathrm{Str}(\mathrm{Crs}_{2r+1}(C_F(c)); \gamma_2) \tag{238}$$

where $r = ||F|| - 1$ and

$$\begin{aligned} \gamma_1 &:= \mathrm{Path}((r,r), ((2r+1)L + r, r)), \\ \gamma_2 &:= \mathrm{Path}(\mathbf{0}, (L,0)). \end{aligned} \tag{239}$$

VII. TOPOLOGICAL STABILIZER GROUPS

This section uncovers the close relationship between $C_{\mathcal{S}}$ and $C_{\mathcal{G}}$. A classification of the topological charges that a topological gauge group may exhibit is given, together with examples that realize each of the different possibilities.

A. Charge groups

Let $\mathcal{S}$ be a TSSG with gauge group $\mathcal{G}$, which is a LPG according to corollary 13. The functions

$$\begin{aligned} \kappa &: C_{\mathcal{G}} \times C_{\mathcal{G}} \longrightarrow \pm 1, \\ \theta &: C_{\mathcal{G}} \qquad\quad\; \longrightarrow \pm 1, \\ \kappa &: C_{\mathcal{G}} \times C_{\mathcal{S}} \longrightarrow \pm 1, \end{aligned} \tag{240}$$

are all defined. Thus the commutation rules of string operators are, for many geometries, dictated by the topology and the charge labels. In this sense, the charges in $C_{\mathcal{G}}$ and $C_{\mathcal{S}}$ are topological.

The charges in $C_{\mathcal{G}}$ and $C_{\mathcal{S}}$ are naturally related. Since $\mathcal{S} \subset \mathcal{G}$, there is an injection morphism

$$\iota : \mathcal{S} \longrightarrow \mathcal{G}. \tag{241}$$

This induces the morphism

$$\begin{aligned} \iota^* \; : \; \Phi^0(\mathcal{G}) &\longrightarrow \Phi^0(\mathcal{S}) \\ \phi &\longmapsto \phi \circ \iota, \end{aligned} \tag{242}$$

which clearly satisfies

$$\iota^*[\mathrm{Com}_{\mathcal{G}}[\mathcal{P}]] \subset \mathrm{Com}_{\mathcal{S}}[\mathcal{P}]. \tag{243}$$

Therefore, ι^* gives rise to a morphism

$$C_\iota : C_{\mathcal{G}} \longrightarrow C_{\mathcal{S}}. \tag{244}$$

Proposition 32 *Let $c, d \in C_{\mathcal{G}}$. Then*

$$\kappa(c,d) = \kappa(c, C_\iota(d)). \tag{245}$$

Proof — By coarse graining, we can always get $\mathcal{S}_g$ and $\mathcal{G}_g$ such that proposition 3 holds for them and for every $s \in \mathcal{S}_g$ there exists $G_s \subset \mathcal{G}_g$ with $s \in \mathrm{Pro}(G_s)$ and $\mathrm{Sup}(G_s) \subset \mathrm{ThkSup}(s)$. Then just notice that for any $c \in C_{\mathcal{G}}$, $L \in \mathbf{Z}$,

$$\begin{aligned} &\mathrm{Crs}_5(\mathrm{Str}(c; \mathrm{Path}((2,2),(5L+2,2)))) \subset \\ &\qquad \subset \mathrm{Str}(\mathrm{Crs}_5(C_\iota(c)); \mathrm{Path}(\mathbf{0},(L,0))). \end{aligned} \tag{246}$$

□

Proposition 33 *Let $c \in C_{\mathcal{G}}$. Then*

$$C_\iota(c) = 1 \quad \Longleftrightarrow \quad \forall d \in C_{\mathcal{G}} \quad \kappa(c,d) = 1. \tag{247}$$

Proof — If $C_\iota(c) = d \neq 1$ for some $d \in C_{\mathcal{S}}$, according to proposition 30 there exists $e \in C_{\mathcal{G}}$ such that $\kappa(e,d) = -1 = \kappa(e,c)$. □

Definition 34 *The TSSGs $\mathcal{S}_i$, $i = 1,2$, have isomorphic topological charges if there exists a group isomorphism*

$$\lambda : C_{\mathcal{G}_1} \longrightarrow C_{\mathcal{G}_2}, \tag{248}$$

such that, when both sides can be applied,

$$\kappa \circ (\lambda_{\mathcal{G}} \times \lambda_{\mathcal{G}}) = \kappa, \qquad \theta \circ \lambda_{\mathcal{G}} = \theta, \tag{249}$$

A TSSG isomorphism F from $\mathcal{S}_1$ to $\mathcal{S}_2$ induces a topological charge isomorphism setting $\lambda = C_F$, with F regarded as an isomorphism from $\mathcal{G}_1$ to $\mathcal{G}_2$.

B. Canonical charge generators

Theorem 35 *Let $\mathcal{S}$ be a TSSG with gauge group $\mathcal{G}$. For some $\alpha, \beta \in \mathbf{N}_0$, $C_{\mathcal{S}}$ and $C_{\mathcal{G}}$ admit independent set of generators*

$$\begin{aligned} C_{\mathcal{G}} &= \langle c_1, \ldots, c_\alpha, d_1, \ldots, d_\alpha, e_1, \ldots, e_\beta \rangle, \\ C_{\mathcal{S}} &= \langle \tilde{c}_1, \ldots, \tilde{c}_\alpha, \tilde{d}_1, \ldots, \tilde{d}_\alpha, \tilde{e}_1, \ldots, \tilde{e}_\beta \rangle, \end{aligned} \tag{250}$$

such that, for $i,j=1,\dots,\alpha$ and $k,l=1,\dots,\beta$,

$$C_\iota(c_i)=\tilde{c}_i,\qquad C_\iota(d_i)=\tilde{d}_i,\qquad C_\iota(e_k)=1, \tag{251}$$

$$\kappa(c_i,c_j)=\kappa(d_i,d_j)=1,\quad \kappa(c_i,d_j)=1-2\delta_{ij}, \tag{252}$$

$$\kappa(c_i,\tilde{e}_k)=\kappa(d_i,\tilde{e}_k)=1,\quad \kappa(e_k,\tilde{e}_l)=1-2\delta_{kl}, \tag{253}$$

$$\theta(c_i)=\theta(d_i)=f_1^{\delta_{i1}},\qquad \theta(e_i)=f_2^{\delta_{k1}}, \tag{254}$$

where $f_1,f_2=\pm1$ only depend on the TSSG, not the choice of generators, and the same is true for α, β.

Proof — We find the generators in three steps.
1) Let K be the kernel of C_ι in $C_{\mathcal{G}}$. Notice that

$$\kappa(e_1,e_2)=\kappa(e_1,C_\iota(e_2))=\kappa(e_1,1)=1 \tag{255}$$

for any $e_1,e_2\in K$. $K\simeq\mathbf{Z}_2^\beta$ for some $\beta\in\mathbf{N}_0$ and it admits a set of independent generators $K=\langle e_1,\dots,e_\beta\rangle$. If $\theta(e_k)=1$ for every $k=1,\dots,n$, we do nothing and set $f_2=1$. In other case, suppose w.l.o.g. that $\theta(e_1)=-1$ and take new generators $\bar{e}_k$ with $\bar{e}_k=e_k$ if $\theta(e_k)=1$ or $k=1$, $\bar{e}_k=e_ke_\beta$ otherwise, remove the bars and set $f_2=-1$. Since $f_2=1$ iff all the elements of K are bosons, f_2 does not depend on the choice of generators.
2) $C_{\mathcal{G}}\simeq\mathbf{Z}_2^{\beta+n}$ for some $n\in\mathbf{N}_0$ and we choose a set of independent generators $\mathcal{G}=\langle e_1,\dots,e_\beta,g_1,\dots,g_n\rangle$. According to proposition 33, for each g_i there exists g_j such that $\kappa(g_i,g_j)=-1$, $i,j=1,\dots,n$. In a similar way that a canonical basis of the Pauli group is obtained, we can obtain from the g_i a set of generators of $C_{\mathcal{G}}$ as in (250) and satisfying (252). If $\theta(c_i)\theta(d_i)=1$ does not hold, we can always find suitable generators. E.g., if $\theta(c_i)=-\theta(d_i)=1$ we set $\bar{c}_i:=c_i,\bar{d}_i:=c_id_i$ and remove the bars. Similarly, suppose that there exist $1\leq i<j\leq n$ with $\theta(c_i)=\theta(c_j)=-1$. Then we can set $\bar{c}_i:=c_ic_j$, $\bar{d}_i:=d_ic_j$, $\bar{c}_j:=c_id_id_j$ and $\bar{d}_j:=c_id_ic_jd_j$ so that $\theta(\bar{c}_i)=\theta(\bar{d}_i)=\theta(\bar{c}_j)=\theta(\bar{d}_j)=1$, and remove the bars. f_1 only depends on the TSG because the total number of bosonic charges in $C_{\mathcal{G}}$ is

$$2^{\alpha+\beta-1}(2^{\alpha+1}+f_1+f_1f_2). \tag{256}$$

3) We define $\tilde{c}_i=C_\iota(c_i)$, $\tilde{d}_i=C_\iota(d_i)$, $i=1,\dots,\alpha$. The set $\{c_1,\dots,c_\alpha,d_1,\dots,d_\alpha\}$ is independet. E.g., $\tilde{c}_i=\tilde{c}_j$ only if $i=j$, because $\kappa(d_i,\tilde{c}_j)=\kappa(d_i,c_j)=1-2\delta_{ij}$. Finally, for each e_k we choose $\tilde{e}_k$ such that $\kappa(e_k,\tilde{e}_k)=-1$, $k=1,\dots,\beta$. If (253) does not hold, we can adjust it easily. E.g., if $\kappa(e_k,\tilde{e}_l)=-1$ for $l>k$, set $\overline{\tilde{e}_l}:=\tilde{e}_k\tilde{e}_l$, and then remove the bars. Or, if $\kappa(c_i,\tilde{e}_k)=-1$, set $\overline{\tilde{e}_k}:=\tilde{e}_k\tilde{d}_i$, and then remove the bars, and so on. □

Definition 36 *The characteristic of a TSSG is given by the numbers $\alpha,\beta\in\mathbf{N}_0$ and $f_1,f_2=\pm1$ in theorem 35, with $f_1=1$ if $\alpha=0$ and $f_2=1$ if $\beta=0$.*

Notice that two TSSGs have isomorphic topological charges iff they have the same characteristic. Some examples follow.

- Empty code, trivial code, trivial subsystem code:
$$\alpha=0,\beta=0,f_1=1,f_2=1. \tag{257}$$
- Toric code:
$$\alpha=1,\beta=0,f_1=1,f_2=1. \tag{258}$$
- Subsystem toric code:
$$\alpha=0,\beta=1,f_1=1,f_2=1. \tag{259}$$
- Topological subsystem color codes:
$$\alpha=1,\beta=0,f_1=-1,f_2=1. \tag{260}$$
- Honeycomb subsystem code:
$$\alpha=0,\beta=1,f_1=1,f_2=-1. \tag{261}$$

These examples and the following observation show that there exist TSSGs of arbitrary characteristic.

Proposition 37 *The composition of two TSSGs with characteristics $\alpha^k,\beta^k,f_1^k,f_2^k$, $k=1,2$, has characteristic*

$$\begin{aligned}
&\alpha=\alpha_1+\alpha_2,\\
&\beta=\beta_1+\beta_2,\\
&f_1=f_1^1f_1^2,\\
&f_2=(1+f_2^1)(1+f_2^2)/2-1.
\end{aligned} \tag{262}$$

C. Characteristic of TSGs

A TSG $\mathcal{S}$ can be regarded as a TSSG with $\beta=0$, so that its characteristic is given by α and f_1. But all known TSGs have $f=1$ (e.g., for toric codes $\alpha=1$ and for color codes $\alpha=2$). Indeed, from the condensed matter perspective the Hamiltonian model related to a TSG is chiral if $f_1=-1$ [33], and this is thought to be incompatible with the fact that the stabilizer generators commute with each other. Therefore, unlike in TSSGs not all characteristics may admit a realization. On the other hand, if we consider only non-chiral TSGs, those with $f_1=1$, there exists TSGs with arbitrary values of α.

VIII. HOMOLOGICAL STRUCTURE

This section discusses how any TSSG can be rearranged in a standard form by means of a framework of string operators. Several examples illustrate the construction. Other concrete examples —that originally motivated the general construction— can be found in [33].

A. Results

For charges as in theorem 35, we use the notation

$$\begin{aligned} K_1 &:= \{1, \dots, \alpha\}, \\ K_2 &:= \{\alpha+1, \dots, 2\alpha\}, \\ K_3 &:= \{2\alpha+1, \dots, 2\alpha+\beta\}, \\ K &:= K_1 \cup K_2 \cup K_3, \end{aligned} \tag{263}$$

$$k^* := \begin{cases} k+\alpha & \text{if } k \in K_1, \\ k-\alpha & \text{if } k \in K_2, \\ k & \text{if } k \in K_3, \end{cases} \tag{264}$$

$$\begin{aligned} &c_k := d_{k-\alpha}, \qquad \tilde{c}_k := \tilde{d}_{k-\alpha}, \qquad k \in K_2, \\ &c_k := e_{k-2\alpha}, \qquad \tilde{c}_k := \tilde{e}_{k-2\alpha}, \qquad k \in K_3. \end{aligned} \tag{265}$$

Proposition 38 *Any TSSG can be coarse grained to a TSSG $\mathcal{S}$, with gauge group $\mathcal{G}$ and LTI set of independent generators $\mathcal{S}_g$, $\mathcal{G}_g$, satisfying the properties in proposition 22 and also the following. There exist*

$$s_k \in \mathcal{S}_g, \qquad g_k \in \mathcal{G}_g - \langle i\mathbf{1}\rangle, \qquad k \in K, \tag{266}$$

with

$$\mathrm{Mor}_{\mathcal{S}_g}(\{s_k\}) \in \tilde{c}_k, \qquad \mathrm{Mor}_{\mathcal{G}_g}(\{g_k\}) \in c_k, \tag{267}$$

$$\mathrm{Sup}(s_k) = \mathrm{Sup}(g_k) = \square^{1,1}_{0,0}, \tag{268}$$

and, for $k \in K_1 \cup K_2$,

$$\mathrm{Mor}_{\mathcal{S}_g}(\{s_k\}) = \iota^*(\mathrm{Mor}_{\mathcal{G}_g}(\{g_k\})). \tag{269}$$

Proof — Coarse graining and choosing $\mathcal{G}'_g, \mathcal{S}'_g$ according to proposition 22 for certain $c'_k, \tilde{c}'_k$, there exist s'_k, g'_k that satisfy all conditions except (269). Let $l \in \mathbf{N}^*$ be the minimal integer such that for any $k \in K_1 \cup K_2$

$$\mathrm{Sup}_{\mathcal{G}'_g}(\iota^*(g_k)) \subset \mathrm{Thk}^l(\square^{1,1}_{0,0}). \tag{270}$$

Now set

$$\begin{aligned} \mathcal{G}_g &:= \mathrm{Crs}_{2l+1}(\mathcal{G}'_g), \\ \mathcal{S}''_g &:= \mathrm{Crs}_{2l+1}(\mathcal{S}'_g), \\ c_k &:= \mathrm{Crs}_{2l+1}(c'_k), \\ \tilde{c}_k &:= \mathrm{Crs}_{2l+1}(\tilde{c}'_k), \end{aligned} \tag{271}$$

and

$$s''_k := \mathrm{Crs}_{2l+1}(T_{l,l}\,(s'_k)), \qquad g_k := \mathrm{Crs}_{2l+1}(T_{l,l}\,(g'_k)), \tag{272}$$

so that all properties but (269) are met and for any $k \in K_1 \cup K_2$ we have

$$\mathrm{Sup}(\iota^*(\mathrm{Mor}_{\mathcal{G}'_g}(g_k)) = \square^{1,1}_{0,0}. \tag{273}$$

For $k \in K_3$ we set $s_k := s''_k$. Let us show that there exist a list of LTI sets of independent generators

$$\mathcal{S}''_g = \mathcal{S}^0_g, \mathcal{S}^1_g, \dots, \mathcal{S}^{2\alpha}_g = \mathcal{S}_g \tag{274}$$

such that for any $k \in K$ and $\mathcal{S}^t_g$ with $k > 2\alpha - t$ there exist a suitable s_k. We construct $\mathcal{S}^t_g$ from $\mathcal{S}^{t-1}_g$ as follows. Set $u := 2\alpha + 1 - t$, $\phi := \iota^*(\mathrm{Mor}_{\mathcal{G}_g}(g_u))$ and choose $\hat{s}_0 \in \overline{\mathrm{Ker}_{\mathcal{S}^{t-1}_g}}(\phi)$ such that $\hat{a}_0 \neq a_k$ for any $k \in K$ with $k > u$. This is always possible because the charge c_u is independent of the charges $c_{u+1}, \dots, c_{2\alpha+\beta}$. Label the elements of $\overline{\mathrm{Ker}_{\mathcal{S}^{t-1}_g}}(\phi) - \{\hat{s}_0\}$ as $\{\hat{s}_i\}_{i=1}^r$ and apply proposition 21 repeatedly to perform the substitutions $\hat{s}_i \to \hat{s}_0\hat{s}_i$, $i = 1, \dots, r$. Letting $\mathcal{S}^t_g$ be the resulting set of generators, we get as needed

$$\mathrm{Mor}_{\mathcal{S}^t_g}(\{s_u\}) = \iota^*(\mathrm{Mor}_{\mathcal{G}_u}(\{g_k\})) \in c_u. \tag{275}$$

Notice that the transformations $\hat{s}_i \to \hat{s}_0\hat{s}_i$ may affect the s_k, $k > u$, that had the required properties in $\mathcal{S}^{t-1}_g$, but does not affect these properties. □

Theorem 39 *Let $\mathcal{S}$ be a TSSG with gauge group $\mathcal{G}$, characteristic α, β, f_1, f_2 and canonical generators (250). There exist*

$$L \in \mathbf{N}^*, \qquad M, N \in \mathbf{N}_0, \tag{276}$$

and mappings

$$\begin{aligned} \epsilon_k, \bar{\epsilon}_l &: \Gamma_{\mathrm{edg}} \cup \Gamma^*_{\mathrm{edg}} \longrightarrow \mathcal{P} \\ \pi_k &: \Gamma_{\mathrm{fc}} \cup \Gamma^*_{\mathrm{fc}} \longrightarrow \mathcal{S}, \\ \bar{\pi}_l &: \Gamma_{\mathrm{fc}} \cup \Gamma^*_{\mathrm{fc}} \longrightarrow \mathcal{G}, \\ \mu_m &: \Sigma \longrightarrow \mathcal{P}, \\ \nu_m &: \Sigma \longrightarrow \mathcal{S}, \\ \xi_n &: \Sigma \longrightarrow \mathcal{G}, \end{aligned} \tag{277}$$

where $k = 1, \dots, \alpha$, $l = 1, \dots, \beta$, $m = 1, \dots, M$, $n = 1, \dots, N$, that satisfy the following properties. We use the notation

$$\begin{aligned} Q_1 &:= \{1, \dots, \alpha\}, \\ Q_2 &:= \{\alpha+1, \dots, \alpha+\beta\}, \\ Q_3 &:= \{\alpha+\beta+1, \dots, \alpha+\beta+M\}, \\ Q &:= Q_1 \cup Q_2 \cup Q_3. \end{aligned} \tag{278}$$

1. *The mappings are translationally invariant,*

   $$T_{iL,jL} \circ \square = \square \circ T_{i,j}\,,$$

   for

   $$\square = \epsilon_k, \pi_k, \bar{\epsilon}_l, \bar{\pi}_l, \mu_m, \nu_m, \xi_n. \tag{279}$$

   In addition

   $$\bar{\epsilon}_l[\Gamma_{\mathrm{edg}}] \subset \mathcal{G}, \qquad \bar{\pi}_l[\Gamma_{\mathrm{fc}}] \subset \mathcal{S}. \tag{280}$$

2. *For any* $d, e \in \Gamma_{\mathrm{edg}}$*,* $k = 1, \dots, \alpha$*,* $l = 1, \dots, \beta$*,*

$$(\epsilon_k(d); \epsilon_k(e)) = \begin{cases} \theta(c_k) & \text{if } \partial\, d \cap \partial\, e \neq \emptyset, \\ 1 & \text{otherwise,} \end{cases}$$
$$(\epsilon_k(d^*); \epsilon_k(e^*)) = \begin{cases} \theta(d_k) & \text{if } \partial\, d^* \cap \partial\, e^* \neq \emptyset, \\ 1 & \text{otherwise,} \end{cases}$$
$$(\bar\epsilon_l(d); \bar\epsilon_l(e)) = \begin{cases} \theta(e_l) & \text{if } \partial\, d \cap \partial\, e \neq \emptyset, \\ 1 & \text{otherwise,} \end{cases}$$
$$(\epsilon_k(d); \epsilon_k(e^*)) = (\bar\epsilon_l(d); \bar\epsilon_l(e^*)) = 1 - 2\delta_{d,e}\,. \tag{281}$$

3. *For any* $\sigma_1, \sigma_2 \in \Sigma$*,* $m = 1, \dots, M$*,*

$$(\mu_m(\sigma_1); \nu_m(\sigma_2)) = 1 - 2\delta_{\sigma_1,\sigma_2},$$
$$(\mu_m(\sigma_1); \mu_m(\sigma_2)) = (\nu_m(\sigma_1); \nu_m(\sigma_2)) = 1. \tag{282}$$

4. *For any* $F \in \mathbf{P}_0(\Gamma_{\mathrm{fc}} \cup \Gamma^*_{\mathrm{fc}})$*,*

$$\mathrm{Pro}(\pi_k[F]) = \mathrm{Pro}(\epsilon_k[\partial F]),$$
$$\mathrm{Pro}(\bar\pi_l[F]) = \mathrm{Pro}(\bar\epsilon_l[\partial F]), \tag{283}$$

where $k = 1, \dots, \alpha$*,* $l = 1, \dots, \beta$*.*

5. *Define the stabilizer and gauge group subsets*

$$\mathcal{S}^k_g := \begin{cases} \pi_k[\Gamma_{\mathrm{fc}}] & \text{if } k \in K_1, \\ \pi_{k-\alpha}[\Gamma^*_{\mathrm{fc}}] & \text{if } k \in K_2, \\ \bar\pi_{k-2\alpha}[\Gamma_{\mathrm{fc}}] & \text{if } k \in K_3, \end{cases}$$
$$\mathcal{G}^k_g := \begin{cases} \pi_k[\Gamma_{\mathrm{fc}}] & \text{if } k \in K_1, \\ \pi_{k-\alpha}[\Gamma^*_{\mathrm{fc}}] & \text{if } k \in K_2, \\ \bar\pi_{k-2\alpha}[\Gamma^*_{\mathrm{fc}}] & \text{if } k \in K_3, \end{cases} \tag{284}$$

The sets

$$\mathcal{S}_g := \bigsqcup_{k\in K} \mathcal{S}^k_g \sqcup \mathcal{S}^0_g, \qquad \mathcal{S}^0_g := \bigsqcup_{m=1}^{M} \nu_m[\Sigma],$$
$$\mathcal{G}_g := \bigsqcup_{k\in K} \mathcal{G}^k_g \sqcup \mathcal{G}^0_g \sqcup \{i\mathbf{1}\}, \qquad \mathcal{G}^0_g := \bigsqcup_{n=1}^{N} \xi_n[\Sigma], \tag{285}$$

are LTI sets of independent generators of $\mathcal{S}$ *and* $\mathcal{G}$*, respectively, with charges*

$$\mathrm{Mor}_{\mathcal{S}_g}(\{s\}) \in \tilde c_{k^*}, \qquad s \in \mathcal{S}^k_g,$$
$$\mathrm{Mor}_{\mathcal{S}_g}(\{s\}) \in \mathrm{Com}_{\mathcal{S}}[\mathcal{P}], \qquad s \in \mathcal{S}^0_g,$$
$$\mathrm{Mor}_{\mathcal{G}_g}(\{g\}) \in c_{k^*}, \qquad s \in \mathcal{G}^k_g,$$
$$\mathrm{Mor}_{\mathcal{G}_g}(\{g\}) \in \mathrm{Com}_{\mathcal{G}}[\mathcal{P}], \qquad g \in \mathcal{G}^0_g. \tag{286}$$

Moreover, for $E \subset \Gamma_{\mathrm{edg}}$*,* $E' \subset \Gamma^*_{\mathrm{edg}}$*,* $\gamma \subset \Sigma$*,*

$$\overline{\mathrm{Ker}_{\mathcal{S}_g}}(\epsilon_k[EE']) = \pi_k[\partial(EE')],$$
$$\overline{\mathrm{Ker}_{\mathcal{S}_g}}(\bar\epsilon_l[E']) = \bar\pi_l[\partial E'],$$
$$\overline{\mathrm{Ker}_{\mathcal{S}_g}}(\mu_m[\gamma]) = \nu_m[\gamma],$$
$$\overline{\mathrm{Ker}_{\mathcal{G}_g}}(\epsilon_k[EE']) = \pi_k[\partial(EE')],$$
$$\overline{\mathrm{Ker}_{\mathcal{G}_g}}(\bar\epsilon_l[E]) = \bar\pi_l[\partial E], \tag{287}$$

where $k = 1, \dots, \alpha$*,* $l = 1, \dots, \beta$*,* $m = 1, \dots, M$*.*

6. $\mathcal{P}/\mathcal{G}$ *has the direct product structure*

$$\frac{\mathcal{P}}{\mathcal{G}} = \frac{\mathcal{P}^1\mathcal{G}}{\mathcal{G}} \times \cdots \times \frac{\mathcal{P}^{\alpha+\beta+M}\mathcal{G}}{\mathcal{G}}, \tag{288}$$

where

$$\mathcal{P}^q := \langle i\mathbf{1}\rangle\langle\mathcal{P}^q_g\rangle, \qquad q \in Q, \tag{289}$$

with

$$\mathcal{P}^q_g := \begin{cases} \epsilon_q[\Gamma_{\mathrm{edg}} \cup \Gamma^*_{\mathrm{edg}}] & \text{if } q \in Q_1, \\ \bar\epsilon_{q-\alpha}[\Gamma^*_{\mathrm{edg}}] & \text{if } q \in Q_2, \\ \mu_{q-\alpha-\beta}[\Sigma] \cup \nu_{q-\alpha-\beta}[\Sigma] & \text{if } q \in Q_3. \end{cases} \tag{290}$$

7. *Let for* $q \in Q_2$

$$\tilde{\mathcal{G}}^q := \langle i\mathbf{1}\rangle\langle\tilde{\mathcal{G}}^q_g\rangle, \qquad \tilde{\mathcal{G}}^q_g := \bar\epsilon_{q-\alpha}[\Gamma_{\mathrm{edg}}]. \tag{291}$$

Then, for $q \neq q'$*,*

$$\mathcal{P}^q \subset \mathcal{Z}_{\mathcal{P}}(\mathcal{P}^{q'}), \qquad q \in Q_1 \cup Q_3, q' \in Q,$$
$$\tilde{\mathcal{G}}^q \subset \mathcal{Z}_{\mathcal{P}}(\mathcal{P}^{q'}), \qquad q \in Q_2, q' \in Q_1 \cup Q_2. \tag{292}$$

8. $\mathcal{S}$ *has the direct product structure*

$$\mathcal{S} = \tilde{\mathcal{S}}^1 \times \cdots \times \tilde{\mathcal{S}}^{\alpha+\beta+M}, \tag{293}$$

where, for $q \in Q$*,*

$$\tilde{\mathcal{S}}^q_g := \begin{cases} \mathcal{S} \cap \mathcal{P}^q = \mathcal{S}^q\mathcal{S}^{q^*} & \text{if } q \in Q_1, \\ \mathcal{S} \cap \tilde{\mathcal{G}}^q = \mathcal{S}^{q+\alpha} & \text{if } q \in Q_2, \\ \mathcal{S} \cap \mathcal{P}^q = \mathcal{S}^{q+\alpha} & \text{if } q \in Q_3. \end{cases} \tag{294}$$

9. *Finally,*

$$\mathrm{Com}_{\mathcal{P}}[\mathrm{Cnstr}_{\mathcal{G}}(\mathcal{P}_g)] = \mathrm{Com}_{\mathcal{P}}[\mathcal{S}] \prod_{q\in Q_1\cup Q_2} \mathrm{Com}_{\mathcal{P}}[Z^q],$$
$$\mathrm{Com}_{\mathcal{P}}[\mathrm{Cnstr}_{\mathcal{G}}(\mathcal{G}_g)] = \mathrm{Com}_{\mathcal{P}}[\mathcal{S}] \prod_{q\in Q_2} \mathrm{Com}_{\mathcal{P}}[Z^q], \tag{295}$$

where

$$Z^q := \begin{cases} \{\epsilon_q[z] \,:\, z \in \Gamma_{\mathrm{cyc}} \cup \Gamma^*_{\mathrm{cyc}}\} & \text{if } q \in Q_1, \\ \{\bar\epsilon_{q-\alpha}[z] \,:\, z \in \Gamma_{\mathrm{cyc}}\} & \text{if } q \in Q_2. \end{cases} \tag{296}$$

Proof — We proceed in several steps, not following the order of the statements in the proposition.

0) We assume that $\mathcal{G} = \langle\mathcal{G}'_g\rangle$ and $\mathcal{S} = \langle\mathcal{S}'_g\rangle$, with $\mathcal{G}'_g$ and $\mathcal{S}'_g$ LTI set of independent generators, have been coarse grained and chosen according to proposition 38. We can assume such a coarse graining w.l.o.g because it can be absorbed in the value of L.

1) We start constructing $\epsilon_k, \pi_k, \bar\epsilon_l, \bar\pi_l$. Let $L = 3(2\alpha + \beta + 1)$ and set for $k \in K$

$$\phi_k := \mathrm{Mor}_{\mathcal{G}'_g}(\{g_k\}),$$
$$\tilde\phi_k := \mathrm{Mor}_{\mathcal{S}'_g}(\{s_k\}). \tag{297}$$

Choose and set, for $k \in K_1 \cup K_2$,

$$\begin{aligned} H^k &\in \mathrm{Str}(\phi_k, T_{L,0}\,(\phi_k)) \subset \mathrm{Str}(\tilde{\phi}^k, T_{L,0}\,(\tilde{\phi}_k)), \\ V^k &\in \mathrm{Str}(\phi_k, T_{0,L}\,(\phi_k)) \subset \mathrm{Str}(\tilde{\phi}_k, T_{0,L}\,(\tilde{\phi}_k)), \\ u_k &:= \theta(c_k)\,(H^k; T_{L,0}\,(H^k)), \\ v_k &:= u_k\,(H^k; V^k), \\ w_k &:= v_k\,(V^k; T_{0,L}\,(V^k)), \end{aligned} \tag{298}$$

and for $k \in K_3$,

$$\begin{aligned} H^k &\in \mathrm{Str}(\phi_k, T_{L,0}\,(\phi_k)), \\ V^k &\in \mathrm{Str}(\phi_k, T_{0,L}\,(\phi_k)), \\ \tilde{H}^k &\in \mathrm{Str}(\tilde{\phi}^k, T_{L,0}\,(\tilde{\phi}_k)), \\ \tilde{V}^k &\in \mathrm{Str}(\tilde{\phi}_k, T_{0,L}\,(\tilde{\phi}_k)), \\ p_k &\in \mathrm{Str}(\iota^*(\phi_k), 1). \end{aligned} \tag{299}$$

Define $\epsilon_k, \bar{\epsilon}_l$ in terms of (279) setting

$$\begin{aligned} \epsilon_k(\mathbf{e}_1) &:= T_{3k,3k}\,\left(s_k^{1-u_k/2} H^k\right), \\ \epsilon_k(\mathbf{e}_2) &:= T_{3k,3k}\,\left(s_k^{1-v_k/2} (T_{0,L}\,(s_k))^{1-w_k/2} V^k\right), \end{aligned} \tag{300}$$

for $k \in K_1$,

$$\begin{aligned} \epsilon_{k-\alpha}(\mathbf{e}_1^*) &:= T_{3k,3k-L}\,\left(s_k^{\frac{1-v_k}{2}} (T_{0,L}\,(s_k))^{\frac{1-w_k}{2}} V^k\right), \\ \epsilon_{k-\alpha}(\mathbf{e}_2^*) &:= T_{3k-L,3k}\,\left(s_k^{\frac{1-u_k}{2}} H^k\right), \end{aligned} \tag{301}$$

for $k \in K_2$, and

$$\begin{aligned} \bar{\epsilon}_{k-2\alpha}(\mathbf{e}_1) &:= \psi_k^1\, T_{3k,3k}\,(p_k H^k T_{L,0}\,(p_k)), \\ \bar{\epsilon}_{k-2\alpha}(\mathbf{e}_2) &:= \psi_k^2\, T_{3k,3k}\,(p_k V^k T_{0,L}\,(p_k)), \\ \bar{\epsilon}_{k-2\alpha}(\mathbf{e}_1^*) &:= T_{(i+1)L-3,jL-3}\,(\tilde{V}^k), \\ \bar{\epsilon}_{k-2\alpha}(\mathbf{e}_2^*) &:= T_{iL-3,(j+1)L-3}\,(\tilde{H}^k), \end{aligned} \tag{302}$$

for $k \in K_3$, where we choose $\psi_k^1, \psi_k^2 \in \langle i\mathbf{1}\rangle$ in order to get elements of $\mathcal{G}$. From the construction, the string commutation rules and the fact that $\mathcal{S} \subset \mathcal{Z}_{\mathcal{P}}(\mathcal{G})$, (281) follows easily. We also get (292) for $q, q' \notin Q_3$.

The previous construction is such that for any $E_1 \subset \Gamma_{\mathrm{edg}}$, $E_2 \subset \Gamma^*_{\mathrm{edg}}$,

$$\begin{aligned} \mathrm{Com}_{\mathcal{G}}(\epsilon_k[E_1 E_2]) = 1 \quad &\Longleftrightarrow \quad \partial E_1 = \partial E_2 = \emptyset, \\ \mathrm{Com}_{\mathcal{G}}(\bar{\epsilon}_k[E_1]) = 1 \quad &\Longleftrightarrow \quad \partial E_1 = \emptyset, \\ \mathrm{Com}_{\mathcal{S}}(\bar{\epsilon}_k[E_2]) = 1 \quad &\Longleftrightarrow \quad \partial E_2 = \emptyset. \end{aligned} \tag{303}$$

In addition, the homology of the plane is trivial and thus

$$\begin{aligned} \partial E_1 = \emptyset \quad &\Longleftrightarrow \quad \exists F_1 \subset \Gamma_{\mathrm{fc}}\,:\, \partial F_1 = E_1, \\ \partial E_2 = \emptyset \quad &\Longleftrightarrow \quad \exists F_2 \subset \Gamma^*_{\mathrm{fc}}\,:\, \partial F_2 = E_2. \end{aligned} \tag{304}$$

Putting this together, (279, 283) become definitions for $\pi_k, \bar{\pi}_l$, possibly up to a phase in the case of gauge elements, which can in that case be chosen arbitrarily.
2) We next construct $\mathcal{S}_g$, $\mathcal{G}_g$. Let for $i, j \in \mathbf{Z}$,

$$\begin{aligned} s^k_{i,j} &:= T_{iL+3k^*,jL+3k^*}\,(s_{k^*}), && k \in K_1 \cup K_2, \\ s^k_{i,j} &:= T_{(i+1)L-3,(j+1)L-3}\,(s_k), && k \in K_3, \\ g^k_{i,j} &:= T_{iL+3k^*,jL+3k^*}\,(g_{k^*}), && k \in K, \end{aligned} \tag{305}$$

and

$$\begin{aligned} E_{i,j} &:= \bigcup_{k \in \mathbf{N}_0} T_{i+k,j}\,(\mathbf{e}_1), \\ E'_{i,j} &:= \bigcup_{k \in \mathbf{N}^*} T_{i+k,j}\,(\mathbf{e}_2^*). \end{aligned} \tag{306}$$

Then for any $k_1 \in K_1$, $k_2 \in K_2$, $k_3 \in K_3$, by construction

$$\begin{aligned} \overline{\mathrm{Ker}_{\mathcal{G}_g}}(\epsilon_{k_1}[E_{i,j}]) = \overline{\mathrm{Ker}_{\mathcal{S}_g}}(\epsilon_{k_1}[E_{i,j}]) &= \{g^{k_1}_{i,j}\} = \{s^{k_1}_{i,j}\}, \\ \overline{\mathrm{Ker}_{\mathcal{G}_g}}(\epsilon_{k_2-\alpha}[E'_{i,j}]) = \overline{\mathrm{Ker}_{\mathcal{S}_g}}(\epsilon_{k_2-\alpha}[E'_{i,j}]) &= \{g^{k_2}_{i,j}\} = \{s^{k_2}_{i,j}\}, \\ \overline{\mathrm{Ker}_{\mathcal{G}_g}}(\bar{\epsilon}_{k_3-2\alpha}[E_{i,j}] T_{Li+3k_3,Lj+3k_3}\,(\{p_k\})) &= \{g^{k_3}_{i,j}\}, \\ \overline{\mathrm{Ker}_{\mathcal{S}_g}}(\bar{\epsilon}_{k_3-2\alpha}[E'_{i,j}]) &= \{s^{k_3}_{i,j}\}. \end{aligned} \tag{307}$$

Set

$$\begin{aligned} S^k &:= \begin{cases} \pi_k(\mathbf{f}^*) & \text{if } k \in K_1, \\ \pi_{k-\alpha}(\mathbf{f}) & \text{if } k \in K_2, \\ \bar{\pi}_{k-2\alpha}(\mathbf{f}) & \text{if } k \in K_3. \end{cases} \\ G^k &:= \begin{cases} \pi_k(\mathbf{f}^*) & \text{if } k \in K_1, \\ \pi_{k-\alpha}(\mathbf{f}) & \text{if } k \in K_2, \\ \bar{\pi}_{k-2\alpha}(\mathbf{f}^*) & \text{if } k \in K_3. \end{cases} \end{aligned} \tag{308}$$

Due to the string commutation rules,

$$\begin{aligned} (\epsilon_{k_1}[E_{i,j}]; G^k) = (\epsilon_{k_1}[E_{i,j}]; S^k) &= 1 - 2\delta_{i,0}\delta_{j,0}, \\ (\epsilon_{k_2-\alpha}[E'_{i,j}]; G^k) = (\epsilon_{k_2-\alpha}[E'_{i,j}]; S^k) &= 1 - 2\delta_{i,0}\delta_{j,0}, \\ (\bar{\epsilon}_{k_3-2\alpha}[E_{i,j}] T_{Li,Lj}\,(\{p_k\}); G^k) &= 1 - 2\delta_{i,0}\delta_{j,0}, \\ (\bar{\epsilon}_{k_3-2\alpha}[E'_{i,j}]; S^k) &= 1 - 2\delta_{i,0}\delta_{j,0}, \end{aligned} \tag{309}$$

It follows that, for $k, k \in K$, $i, j \in \mathbf{Z}$,

$$\begin{aligned} s^k_{i,j} \in \hat{S}^{k'} &\iff i = j = k - k' = 0, \\ g^k_{i,j} \in \hat{G}^{k'} &\iff i = j = k - k' = 0. \end{aligned} \tag{310}$$

where $\hat{S}^k \in \mathbf{P}_0(\mathcal{S}'_g)$, $\hat{G}^k \in \mathbf{P}_0(\mathcal{G}'_g)$ are such that $S^k \in \mathrm{Pro}(\hat{S}^k)$, $G^k \in \mathrm{Pro}(\hat{G}^k)$. Thus we can define new LTI sets of independent generators

$$\begin{aligned} \mathcal{S}_g &:= \mathcal{S}'_g \cup T^{(L)}\,(\{S^k\}) - T^{(L)}\,(\{s^k_{0,0}\}), \\ \mathcal{G}''_g &:= \mathcal{G}'_g \cup T^{(L)}\,(\{G^k\}) - T^{(L)}\,(\{g^k_{0,0}\}). \end{aligned} \tag{311}$$

With these definitions, the charges of S^k, G^k are according to proposition 21,

$$\begin{aligned}\mathrm{Mor}_{\mathcal{S}_g}(\{S^k\}) &\in \tilde{c}_{k^*},\\ \mathrm{Mor}_{\mathcal{G}''_g}(\{G^k\}) &\in c_{k^*}.\end{aligned} \tag{312}$$

Let us show that the rest of generators have trivial charge. Since

$$\partial\Gamma_{\mathrm{fc}} = \partial\Gamma^*_{\mathrm{fc}} = \emptyset, \tag{313}$$

for any $k \in K$ we have

$$\begin{aligned}\mathcal{S}_g^k &= T^{(L)}\left(\{S^k\}\right) \in \mathrm{Cnstr}(\mathcal{S}_g),\\ \mathcal{G}_g^k &= T^{(L)}\left(\{G^k\}\right) \in \mathrm{Cnstr}(\mathcal{G}''_g).\end{aligned} \tag{314}$$

Together with proposition 25, this shows that the elements of the sets

$$\mathcal{S}_g^0 := \mathcal{S}_g - \bigcup_{k\in K} \mathcal{S}_g^k, \qquad \mathcal{G}_g^{0\prime\prime} := (\mathcal{G}''_g - \bigcup_{k\in K} \mathcal{G}_g^k) - \langle i\mathbf{1}\rangle, \tag{315}$$

have trivial charge in $\mathcal{S}_g$ and $\mathcal{G}''_g$, respectively.

For $k \in K_3$ and some $n_k \in \mathbf{N}_0$, let

$$\{g_j^k\}_{j=1}^{n_k} = \overline{\mathrm{Ker}}_{\mathcal{G}''_g}(T_{3k,3k}\left(p_k\right)) \subset \mathcal{G}_g^{0\prime\prime}, \tag{316}$$

and set

$$\begin{aligned}\mathcal{G}_g^0 := \mathcal{G}_g^{0\prime\prime} \cup \bigcup_{k\in K_3} T^{(L)}\left(\{G^k g_j^k\}_{j=1}^{n_k}\right) -\\ - \bigcup_{k\in K_3} T^{(L)}\left(\{g_j^k\}_{j=1}^{n_k}\right).\end{aligned} \tag{317}$$

Then (286, 287) are direct consequences of the construction, using proposition 21 again.

We have to define ν_m and ξ_n so that (285) and (315, 317) agree. Given $s \in \mathcal{S}_g^0$, let $\sigma_s \in \Sigma$ be the minimal element, in lexicographical ordering, of $\mathrm{Sup}(s)$. Then for any $i, j \in \mathbf{Z}$

$$\sigma_{T_{iL,jL}\left(s\right)} = T_{iL,jL}\left(\sigma_s\right). \tag{318}$$

For some $M \in \mathbf{N}_0$, let us label

$$\{s_0^m\}_{m=1}^M = \{s \in \mathcal{S}_g^0 \, : \, \sigma_s \in \square_{0,0}^{L,L}\}, \tag{319}$$

and, imposing translational invariance, define ν_m setting

$$\nu_m(\mathbf{0}) = s_0^m. \tag{320}$$

The definition of ξ_n is entirely analogous.

3) Set $\mathcal{S}^k := \langle\mathcal{S}_g^k\rangle$, $\mathcal{G}^k := \langle\mathcal{G}_g^k\rangle$, $k \in K$. We next show that for any

$$p \in \mathcal{Z}_{\mathcal{P}}(\prod_{k\in K} \mathcal{S}^k), \tag{321}$$

there exists $g \in \mathcal{G}$ such that

$$pg \in \mathcal{Z}_{\mathcal{P}}(\prod_{q\in Q_1\cup Q_2} \mathcal{P}^q). \tag{322}$$

For each $k \in K_3$, let $F_k \in \mathbf{P}_0(\Gamma^*_{\mathrm{fc}})$ be such that

$$\mathrm{Com}_{\mathcal{G}^k}(p) = \mathrm{Mor}_{\mathcal{G}_g^k}(\bar{\pi}_{k-2\alpha}[F_k]). \tag{323}$$

Since $\mathcal{G}_g^k$ is a constraint, $|F_k|$ is even and thus there exists $E_k \in \mathbf{P}_0(\Gamma_{\mathrm{edg}})$ with $\partial E_k = F_k$. Choosing

$$g^k \in \mathrm{Pro}(\bar{\epsilon}_{k-2\alpha}[E_k]) \subset \mathcal{G} \tag{324}$$

we get

$$p' := p \prod_{k\in K_3} g^k \in \mathcal{Z}_{\mathcal{P}}(\prod_{k\in K} \mathcal{S}^k\mathcal{G}^k). \tag{325}$$

Let for $q_1 \in Q_1$, $q_2 \in Q_2$,

$$\begin{aligned}E_{q_1} &:= \{e \in \Gamma_{\mathrm{edg}} \cup \Gamma^*_{\mathrm{edg}} \, : \, (p'; \epsilon_{q_1}(e^*)) = -1\},\\ \bar{E}_{q_2} &:= \{e \in \Gamma_{\mathrm{edg}} \, : \, (p'; \bar{\epsilon}_{q_2-\alpha}(e^*)) = -1\}.\end{aligned} \tag{326}$$

Due to (325) $\partial E_{q_i} = \emptyset$, $i = 1, 2$, and thus there exist $F_{q_1} \in \mathbf{P}_0(\Gamma_{\mathrm{fc}} \cup \Gamma^*_{\mathrm{fc}})$, $F_{q_2} \in \mathbf{P}_0(\Gamma_{\mathrm{fc}})$,with $E_{q_i} = \partial F_{q_i}$. Then we take

$$g \propto \mathrm{Pro}\left(\prod_{q_1\in Q_1} \pi_{q_1}[F_{q_1}] \prod_{q_2\in Q_2} \bar{\pi}_{q_2}[\bar{F}_{q_2}]\right) \prod_{k\in K_3} g^k. \tag{327}$$

4) Next we construct the functions μ_m and prove the direct product structure of $\mathcal{S}$. Due to the construction in (327), for each m we can choose

$$\overline{s_0^m} \in \mathcal{Z}_{\mathcal{P}}(\prod_{q\in Q_1\cup Q_2} \mathcal{P}^q) \tag{328}$$

with

$$\mathrm{Com}_{\mathcal{S}}(\overline{s_0^m}) = \mathrm{Mor}_{\mathcal{S}_g}(s_0^m). \tag{329}$$

Extend this choice to any $s \in \mathcal{S}_0^g$ imposing translational invariance, that is,

$$\overline{T_{iL,jL}\left(s\right)} = T_{iL,jL}\left(\bar{s}\right). \tag{330}$$

Consider any translationally invariant ordering in $\mathcal{S}_g^0$, that is, such that for any $r, s \in \mathcal{S}_g^0$ and $i, j \in \mathbf{Z}$,

$$r < s \iff T_{iL,jL}\left(s\right) < T_{iL,jL}\left(r\right). \tag{331}$$

For $s \in \mathcal{S}_g^0$, let

$$\epsilon_s := \{r \in \mathcal{S}_g^0 \, : \, r < s \text{ and } (\bar{r}; \bar{s}) = -1\}, \tag{332}$$

observing that $|\epsilon_s| < \infty$. Also, let

$$\hat{s} := \bar{s} \prod_{r\in\epsilon_s} r, \tag{333}$$

which is a translationally invariant definition. Then for any $r, s \in \mathcal{S}_g^0$ we have

$$(r; s) = 1, \quad (\hat{r}; \hat{s}) = 1, \quad (r; \hat{s}) = 1 - 2\delta_{r,s}. \tag{334}$$

Imposing translational invariance, we define μ_m setting

$$\mu_m(\mathbf{0}) = \widehat{s_0^m}. \tag{335}$$

This construction ensures (292) and makes the third equation in (294) trivial. The other two follow from $\mathcal{S} \propto \mathcal{Z}_\mathcal{P}(\mathcal{G})$ and (283, 287, 304). Then the direct product structure (293) of $\mathcal{S}$ follows from the structure of $\mathcal{S}_g$.

5) To prove the direct product structure of $\mathcal{P}/\mathcal{G}$, let $p \in \mathcal{P}$ and

$$\begin{aligned} F_{q_1} &\in \mathbf{P}_0(\Gamma_{\text{fc}} \cup \Gamma^*_{\text{fc}}), && q_1 \in Q_1,\\ F_{q_2} &\in \mathbf{P}_0(\Gamma_{\text{fc}}), && q_2 \in Q_2,\\ \gamma_{q_3} &\in \mathbf{P}_0(\Sigma), && q_3 \in Q_3, \end{aligned} \tag{336}$$

be defined by

$$\begin{aligned} \text{Com}_{\tilde{\mathcal{S}}^{q_1}}(p) &= \text{Mor}_{\mathcal{S}_g^{q_1} \cup \mathcal{S}_g^{q_1^*}}(\pi_{q_1}[F_{q_1}]),\\ \text{Com}_{\tilde{\mathcal{S}}^{q_2}}(p) &= \text{Mor}_{\mathcal{S}_g^{q_2+\alpha}}(\pi_{q_2-\alpha}[F_{q_2}]),\\ \text{Com}_{\tilde{\mathcal{S}}^{q_3}}(p) &= \text{Mor}_{\mathcal{S}_g^{q_3+\alpha}}(\nu_{q_3-\alpha-\beta}[\gamma_{q_3}]). \end{aligned} \tag{337}$$

Reasoning as in (324), there exist $E_{q_1} \in \mathbf{P}_0(\Gamma_{\text{edg}} \cup \Gamma^*_{\text{edg}})$, $E_{q_2} \in \mathbf{P}_0(\Gamma^*_{\text{edg}})$ such that

$$\begin{aligned} p_{q_1} &:= \epsilon_{q_1}[E_{q_1}], && q_1 \in Q_1,\\ p_{q_2} &:= \bar{\epsilon}_{q_2-\alpha}[E_{q_2}], && q_2 \in Q_2,\\ p_{q_3} &:= \mu_{q_3-\alpha-\beta}[\gamma_{q_3}], && q_3 \in Q_3, \end{aligned} \tag{338}$$

satisfy

$$\text{Com}_{\tilde{\mathcal{S}}^q}(pp_q) = 1, \qquad q \in Q. \tag{339}$$

Thus, using (292),

$$p\,\mathcal{G} = \prod_{q \in Q} p_q\,\mathcal{G}. \tag{340}$$

If $\bar{p}_q \in \mathcal{P}^q$ give an alternative decomposition, from $p^2\,\mathcal{G} = \mathcal{G} = \mathcal{Z}_\mathcal{P}(\mathcal{S})$ we get, using (287),

$$p_q \bar{p}_q \in \mathcal{P}^q \cap \mathcal{Z}_\mathcal{P}(\mathcal{S}_q) \subset \mathcal{G}. \tag{341}$$

Thus, $p_q\,\mathcal{G} = \bar{p}_q\,\mathcal{G}$.

6) To prove the first equation in (295), take any $R \in \text{Cnstr}_\mathcal{G}(\mathcal{P}_g)$. Let for $q \in Q_2$,

$$E_q := \{e \in \Gamma_{\text{edg}} \,:\, (R; \bar{\epsilon}_{q-\alpha}(e^*)) = -1\}. \tag{342}$$

Since $\text{Com}_{\mathcal{G}^q}(R) = 1$, $\partial E_q = \emptyset$ and thus $z_q := \epsilon_{q-\alpha}[E_q] \in Z^q$ satisfies

$$\text{Com}_{\mathcal{P}^q}(z_q) = \text{Com}_{\mathcal{P}^q}(R). \tag{343}$$

Set $z := \prod_{q \in Q_2} z_q$ and notice that $\text{Com}_\mathcal{G}(z) = 1$. According to (288) there exist $R_q \subset \mathcal{P}_g^q$, $q \in Q$, and $R_0 \subset \mathcal{G}_g^0$, such that, setting $Q_0 := Q \cup \{0\}$,

$$\text{Com}_\mathcal{P}(Rz) = \text{Com}_\mathcal{P}(\prod_{q \in Q_0} R_q). \tag{344}$$

In particular, $\text{Com}_\mathcal{G}(\prod_{q \in Q_0} R_q) = 1$, which implies, using (287, 281), that $R_q \in Z^q$ for $q \in Q_1$, $R_q = \emptyset$ for $q \in Q_2$ and $R_q \subset \mathcal{S}_g^q$ for $q \in Q_3$. It also follows from (288), together with (287), that for each $g \in \mathcal{G}_g^0$ we can choose $\hat{g} \in \mathcal{G}^0 \prod_{q \in Q_2} \mathcal{P}^q$ such that

$$\text{Com}_\mathcal{G}(\hat{g}) = \text{Mor}_{\mathcal{G}_g^0}(\{g\}). \tag{345}$$

Then, using the fact that $R_q = \emptyset$ for $q \in Q_2 \cup Q_3$,

$$(\hat{g}; R_0) = (\hat{g}; \prod_{q \in Q_0} R_q) = (\hat{g}; Rz) = 1. \tag{346}$$

Thus, $R_0 = \emptyset$ and the result follows noting that for $q \in Q_1 \cup Q_2$ we have $\text{Com}_\mathcal{G}[Z^q] = \{1\}$.

The proof of the second equation in (295) is analogous, but $R \in \text{Cnstr}_\mathcal{G}(\mathcal{G}_g)$ and, due to (285), $R_q \subset \mathcal{S}_g^q$ for $q \in Q_1$, $R_q \subset \mathcal{G}_g^q$ for $q \in Q_2$ and there is no need for any R_q with $q \in Q_3$. ☐

Corolary 40 *Let $\mathcal{S}$ be a TSG and consider the construction in theorem 39 for $\mathcal{G} = \mathcal{S}$. Then*

$$\mathcal{P}^q = \mathcal{Z}_\mathcal{P}(\overline{\mathcal{P}^q}), \qquad q \in Q, \tag{347}$$

and $\mathcal{P}/\langle i\mathbf{1}\rangle$ admits the direct product structure

$$\frac{\mathcal{P}}{\langle i\mathbf{1}\rangle} = \frac{\mathcal{P}^1}{\langle i\mathbf{1}\rangle} \times \cdots \times \frac{\mathcal{P}^{\alpha+M}}{\langle i\mathbf{1}\rangle}. \tag{348}$$

B. Examples

Trivial code — $L = M = 1$, $N = 0$ and

$$\mu_1(\mathbf{0}) := X_{0,0}, \qquad \nu_1(\mathbf{0}) := Z_{0,0}. \tag{349}$$

Subsystem trivial code — $L = 1$, $M = N = 0$ and

$$\xi_1(\mathbf{0}) := X_{0,0}, \qquad \xi_2(\mathbf{0}) := Z_{0,0}, \tag{350}$$

Toric code — $L = 1$, $M = N = 0$ and

$$\begin{aligned} \epsilon_1(\mathbf{e}_1) &:= Z^h_{0,0}, & \epsilon_1(\mathbf{e}_1^*) &:= X^h_{0,0},\\ \epsilon_1(\mathbf{e}_2) &:= Z^v_{0,0}, & \epsilon_1(\mathbf{e}_2^*) &:= X^v_{0,0},\\ \pi_1(\mathbf{f}) &:= S^Z, & \pi_1(\mathbf{f}^*) &:= S^X, \end{aligned} \tag{351}$$

Subsystem toric code — $L = 1$, $M = 0$, $N = 2$,

$$\begin{aligned} \bar{\epsilon}_1(\mathbf{e}_1) &:= Z^h_{0,0}, & \bar{\epsilon}_1(\mathbf{e}_1^*) &:= X^h_{0,0},\\ \bar{\epsilon}_1(\mathbf{e}_2) &:= Z^v_{0,0}, & \bar{\epsilon}_1(\mathbf{e}_2^*) &:= X^v_{0,0},\\ \bar{\pi}_1(\mathbf{f}) &:= S^Z, & \bar{\pi}_1(\mathbf{f}^*) &:= S^X,\\ \xi_1(\mathbf{0}) &:= Z^h_{0,0}, & \xi_2(\mathbf{0}) &:= Z^v_{0,0}. \end{aligned} \tag{352}$$

Honeycomb subsystem code — $L = 1$, $M = 0$,

$$\begin{aligned} \bar{\epsilon}_1(\mathbf{e}_1) &:= T_{1,0}\,(G^X)\,G^Y, & \bar{\epsilon}_1(\mathbf{e}_1^*) &:= Y^1_{1,0}, \\ \bar{\epsilon}_1(\mathbf{e}_2) &:= T_{0,1}\,(G^X)\,G^Z, & \bar{\epsilon}_1(\mathbf{e}_2^*) &:= Z^1_{0,1}, \\ \bar{\pi}_1(\mathbf{f}) &:= S^{\hexagon}, & \bar{\pi}_1(\mathbf{f}^*) &:= G^X G^Y G^Z, \\ \xi_1(\mathbf{0}) &:= G^X G^Y, & \xi_2(\mathbf{0}) &:= G^Y G^Z. \end{aligned} \tag{353}$$

Topological subsystem color code — $L = 3$ due to the $\phi \simeq T_{iL,jL}\,(\phi)$ condition, which gives $M = 16$, $N = 88$. Due to the large value of M and N, we omit the description of μ_i, ν_i, ξ_i. The rest of the construction is given by

$$\begin{aligned} \epsilon_1(\mathbf{e}_1) &:= Z^2_{1,0}X^3_{1,0}Y^1_{1,0}X^5_{1,0}X^2_{2,0}Y^4_{2,0}X^6_{2,0}\cdot \\ &\quad\cdot Z^5_{2,0}Y^4_{3,0}Y^2_{3,0}X^6_{3,0}X^3_{3,0}Y^5_{3,0}Y^1_{3,0}, \\ \epsilon_1(\mathbf{e}_2) &:= Z^1_{0,0}Z^6_{0,1}Y^5_{0,1}Y^3_{0,1}X^1_{0,1}X^4_{0,2}Y^6_{0,2}Y^2_{0,2}\cdot \\ &\quad\cdot Z^3_{0,2}X^4_{0,3}Y^2_{0,3}X^6_{0,3}X^3_{0,3}Y^5_{0,3}Y^1_{0,3}, \\ \pi_1(\mathbf{f}) &:= S^1_{0,0}S^3_{1,0}S^2_{2,0}S^2_{0,1}S^1_{1,1}S^3_{1,2}S^3_{2,0}S^2_{2,1}S^1_{2,2}, \\ \epsilon_1(\mathbf{e}_1^*) &:= T_{2,-2}\,(\epsilon_1(\mathbf{e}_2)), \\ \epsilon_1(\mathbf{e}_2^*) &:= T_{-1,1}\,(\epsilon_1(\mathbf{e}_1)), \\ \pi_1(\mathbf{f}^*) &:= T_{-1,-2}\,(\pi_1(\mathbf{f})), \\ S^0_g &:= \mathcal{S}^{\mathrm{col}}_g \cup T^{(3)}\,(\{S^3_{0,0}, S^2_{2,0}S^1_{2,1}, S^3_{2,1}, S^2_{0,2}S^1_{3,0}\}) - \\ &\quad - T^{(3)}\,(\{S^1_{0,0}, S^2_{0,0}, S^2_{2,0}, S^1_{2,1}, S^2_{2,1}, S^2_{0,2}\}), \\ G^0_g &:= \mathcal{G}^{\mathrm{col}}_g - T^{(3)}\,(\{G^1, T_{2,1}\,(G^1)\}). \end{aligned} \tag{354}$$

IX. LOCAL EQUIVALENCE OF TSGS

The general structure of TSGs given in corollary 40 paves the way to show the equivalence, under a suitable class of local transformations, of TSGs with the same topological content.

Definition 41 *Two TSGs $\mathcal{S}_1, \mathcal{S}_2$ are equivalent if there exists a TSG isomorphism*

$$\mathrm{Crs}_{l_1}(\mathcal{S}_1) \otimes \mathcal{S}_{\mathrm{T}}^{\otimes k_1} \xrightarrow{F} \mathrm{Crs}_{l_2}(\mathcal{S}_2) \otimes \mathcal{S}_{\mathrm{T}}^{\otimes k_2}, \tag{355}$$

for some $k_1, k_2 \in \mathbf{N}_0$ and $l_1, l_2 \in \mathbf{N}^*$.

To check that this actually gives an equivalence $\sim$, let $\mathcal{S}_1 \sim \mathcal{S}_2$ and $\mathcal{S}_2 \sim \mathcal{S}_3$ and let us show that $\mathcal{S}_1 \sim \mathcal{S}_3$. There exist local stabilizer isomorphisms

$$\begin{aligned} \mathrm{Crs}_{l_1}(\mathcal{S}_1) \otimes \mathcal{S}_{\mathrm{T}}^{\otimes k_1} &\xrightarrow{F} \mathrm{Crs}_{l_2}(\mathcal{S}_2) \otimes \mathcal{S}_{\mathrm{T}}^{\otimes k_2}, \\ \mathrm{Crs}_{l_2'}(\mathcal{S}_2) \otimes \mathcal{S}_{\mathrm{T}}^{\otimes k_2'} &\xrightarrow{G} \mathrm{Crs}_{l_3}(\mathcal{S}_3) \otimes \mathcal{S}_{\mathrm{T}}^{\otimes k_3}, \end{aligned} \tag{356}$$

for some $k_1, k_2, k_2', k_3 \in \mathbf{N}_0$, and $l_1, l_2, l_2', l_3 \in \mathbf{N}^*$. Then there exists a TSG isomorphism

$$\mathrm{Crs}_{l_1'}(\mathcal{S}_1) \otimes \mathcal{S}_{\mathrm{T}}^{\otimes k_1'} \xrightarrow{H} \mathrm{Crs}_{l_3'}(\mathcal{S}_3) \otimes \mathcal{S}_{\mathrm{T}}^{\otimes k_3'}, \tag{357}$$

where

$$\begin{aligned} l_1' &:= l_1 l_2', & l_3' &:= l_3 l_2, \\ k_1' &:= k_1 l_2' + k_2' l_2, & k_3' &:= k_3 l_2 + k_2 l_2'. \end{aligned} \tag{358}$$

In particular,

$$H = (\mathrm{Crs}_{l_2}(G) \otimes 1_{\mathrm{T}}^{\otimes k_2 l_2'}) \circ (\mathrm{Crs}_{l_2'}(F) \otimes 1_{\mathrm{T}}^{\otimes k_2' l_2}) \tag{359}$$

where 1_{T} is the identity morphism

$$\mathcal{S}_{\mathrm{T}} \xrightarrow{1_{\mathrm{T}}} \mathcal{S}_{\mathrm{T}}. \tag{360}$$

Theorem 42 *Two TSGs are equivalent if and only if they have isomorphic topological charges.*

Proof — The only if direction follows from previous discussions. Thus, we have to show that if $\mathcal{S}_1 \subset \mathcal{P}_1$ and $\mathcal{S}_2 \subset \mathcal{P}_2$ have the same characteristic α, f_1, there exists a TSG isomorphism F as in (355).

We use the constructions of theorem 39 and corollary 40, applied both to $\mathcal{S}_1$ and $\mathcal{S}_2$. For $x = 1, 2$, we let $L_x \in \mathbf{N}^*$, $M_x \in \mathbf{N}_0$ be the constants appearing in these propositions, and we label the mappings as

$$\begin{aligned} \epsilon^x_k &: \Gamma_{\mathrm{edg}} \cup \Gamma^*_{\mathrm{edg}} && \longrightarrow \mathcal{P}_x \\ \pi^x_k &: \Gamma_{\mathrm{fc}} \cup \Gamma^*_{\mathrm{fc}} && \longrightarrow \mathcal{S}_x, \\ \mu^x_m &: \Sigma && \longrightarrow \mathcal{P}_x, \\ \nu^x_m &: \Sigma && \longrightarrow \mathcal{S}_x, \end{aligned} \tag{361}$$

where $k = 1, \ldots, \alpha$, $m = 1, \ldots, M_x$. Let H^x_k, $\tilde{H}^x_k$, V^x_k, $\tilde{V}^x_k$, S^x_k, $\tilde{S}^x_k$ be self-adjoint operators such that

$$\begin{aligned} H^x_k &\propto \epsilon^x_k(\mathbf{e}_1), \\ V^x_k &\propto \epsilon^x_k(\mathbf{e}_2), \\ \tilde{H}^x_k &\propto \epsilon^x_k(\mathbf{e}_1^*), \\ \tilde{V}^x_k &\propto \epsilon^x_k(\mathbf{e}_2^*), \\ \tilde{S}^x_m &\propto \mu^x_m(\mathbf{0}), \\ S^x_m &= \mu^x_m(\mathbf{0}). \end{aligned} \tag{362}$$

Define for $k = 1, \ldots, \alpha$,

$$\begin{aligned} s^k_x &:= \pi^x_k(\mathbf{f})\, H^k_x\, T_{0,L_x}\,(H^k_x)\, V^k_x\, T_{L_x,0}\,(V^k_x) = \pm 1, \\ \tilde{s}^k_x &:= \pi^x_k(\mathbf{f}^*)\, \tilde{H}^k_x\, T_{0,-L_x}\,(\tilde{H}^k_x)\, \tilde{V}^k_x\, T_{-L_x,0}\,(\tilde{V}^k_x) = \pm 1. \end{aligned} \tag{363}$$

We set for $i, j \in \mathbf{Z}$

$$\begin{aligned} H^k_{x,i,j} &:= (s^k_x)^i\, T_{iL_x,jL_x}\,(H^k_x), \\ \tilde{H}^k_{x,i,j} &:= (\tilde{s}^k_x)^i\, T_{iL_x,jL_x}\,(\tilde{H}^k_x). \end{aligned} \tag{364}$$

For $x = 1, 2$, we consider M_x copies of $\mathcal{P}_{\mathrm{T}}$ and label single qubit pauli operators as $X^m_{x,i,j}$, $Z^m_{x,i,j}$, with $m = 1, \ldots, M_x$ labeling the copy and $i, j \in \mathbf{Z}$ labeling the site. Then we define a TSG isomorphism of step 2

$$\mathrm{Crs}_{L_1}(\mathcal{S}_1) \otimes \mathcal{S}_{\mathrm{T}}^{M_2} \xrightarrow{F} \mathrm{Crs}_{L_2}(\mathcal{S}_2) \otimes \mathcal{S}_{\mathrm{T}}^{M_1} \tag{365}$$

setting for $i,j \in \mathbf{Z}$, $k = 1,\dots,\alpha$, $m_1 = 1,\dots,M_1$, $m_2 = 1,\dots,M_2$,

$$\begin{aligned} F(H^k_{1,i,j}) &= H^k_{2,i,j}, & F(V^k_{1,i,j}) &= V^k_{2,i,j}, \\ F(\tilde H^k_{1,i,j}) &= \tilde H^k_{2,i,j}, & F(\tilde V^k_{1,i,j}) &= \tilde V^k_{2,i,j}, \\ F(\nu^1_{m_1}((i,j))) &= Z^{m_1}_{1,i,j}, & F(\mu^1_{m_1}((i,j))) &= X^{m_1}_{1,i,j}, \\ F(Z^{m_2}_{2,i,j}) &= \nu^2_{m_2}((i,j)), & F(X^{m_2}_{2,i,j}) &= \mu^2_{m_2}((i,j)). \end{aligned} \tag{366}$$

□

X. CODES IN THE TORUS

With a bit of care, the structure of TSSGs unveiled in theorem 39 can be used to understand their counterparts in finite lattices, that is, TSCs. This section sketches how to do this for lattices with periodic boundary conditions, so that the topology is that of a torus. We take the periodicity of the lattice to be a multiple of the constant L appearing in the structure theorem. Other periodicities would not change the local structure, but could give rise to global defects (twist lines). Another variation would be to consider a system with boundaries rather than a closed surface, but this is beyond the scope of this paper.

A key point is that we can *reconstruct* the code on the finite lattice by reinterpreting in the natural way the mappings (277). Namely, if the lattice has dimensions $xL \times yL$ for some $x, y \in \mathbf{Z}$, we take Γ to be a periodic square lattice of dimensions $x \times y$, rather than an infinite lattice. The only constraint that we impose is that x and y are not too small. In particular, if m is the maximal range among the image elements of the original mappings (277), we require that $xL, yL \geq 2m$. This ensures that all the properties involving local commutation relations remain true, for example (292, 281, 282, 287). The relations (283) are still true too, and from (288) it follows easily —consider single qubit operators— that

$$\mathcal{P} = \mathcal{P}^1 \cdots \mathcal{P}^{\alpha+\beta+M}\mathcal{G}. \tag{367}$$

For convenience, for $q \in Q_2$ set $\epsilon_{\alpha+q} := \bar\epsilon_q$, $\pi_{\alpha+q} := \bar\pi_q$. We define for $q \in Q$ the cycle and boundary groups of operators

$$\begin{aligned} \mathcal{Z}_q &:= \bigcup\{\mathrm{Pro}(\epsilon_q[z]) \,:\, z \in \Gamma_{\mathrm{cyc}}\}, \\ \mathcal{Z}^*_q &:= \bigcup\{\mathrm{Pro}(\epsilon_q[z]) \,:\, z \in \Gamma^*_{\mathrm{cyc}}\}, \\ \mathcal{B}_q &:= \bigcup\{\mathrm{Pro}(\pi_q[F]) \,:\, F \subset \Gamma_{\mathrm{fc}}\}, \\ \mathcal{B}^*_q &:= \bigcup\{\mathrm{Pro}(\pi_q[F]) \,:\, F \subset \Gamma^*_{\mathrm{fc}}\}, \end{aligned} \tag{368}$$

and choose self-adjoint cycle operators

$$\begin{aligned} z_{1,q} &\in \mathrm{Pro}(\epsilon_q(\mathbf{z}_1)), & z_{2,q} &\in \mathrm{Pro}(\epsilon_q(\mathbf{z}_2)), \\ z^*_{1,q} &\in \mathrm{Pro}(\epsilon_q(\mathbf{z}^*_2)), & z^*_{2,q} &\in \mathrm{Pro}(\epsilon_q(\mathbf{z}^*_1)), \end{aligned} \tag{369}$$

which, due to (281), satisfy the relations, for $i,j = 1,2$,

$$\begin{aligned} (z_{i,q}; z^*_{j,q'}) &= 1 - 2\delta_{ij}\delta_{qq'}, & q,q' &\in Q, \\ (z_{i,q}; z_{j,q'}) &= 1, & q,q' &\in Q, \\ (z^*_{i,q}; z^*_{j,q'}) &= 1, & q \in Q&, q' \in Q_1. \end{aligned} \tag{370}$$

According to the homology of the torus

$$\begin{aligned} \mathcal{Z}_q &= \langle z_{1,q}, z_{2,q}\rangle \mathcal{B}_q, \\ \mathcal{Z}^*_q &= \langle z^*_{1,q}, z^*_{2,q}\rangle \mathcal{B}^*_q, \end{aligned} \tag{371}$$

and notice that

$$\begin{aligned} \mathcal{B}_q &\propto \mathcal{S}^q, \\ \mathcal{B}^*_q &\propto \begin{cases} \mathcal{S}^{q^*} & \text{if } q \in Q_1, \\ \mathcal{G}^{q+\alpha} & \text{if } q \in Q_2. \end{cases} \end{aligned} \tag{372}$$

In the finite lattice $\mathcal{G}_g$ and $\mathcal{S}_g$ are not necessarily independent sets of generators anymore. Indeed, the global constraints become now the relations:

$$\prod_{g \in \mathcal{G}^k_g} g \propto 1, \qquad \prod_{s \in \mathcal{S}^k_g} s \propto 1. \tag{373}$$

No other such independent relations can exist among subsets of $\mathcal{G} = \langle \mathcal{G}_g\rangle$ or $\mathcal{S} := \langle \mathcal{S}_g\rangle$. Also, $\mathcal{S}$ is clearly abelian and its elements self-adjoint, so that

$$\prod_{s \in \mathcal{S}^k_g} s = \pm 1 \tag{374}$$

and, if necessary, we can change the sign of a stabilizer generator to make the product positive. This makes $\mathcal{S}$ a stabilizer group.

Similarly, in the finite lattice the second equation in (295) becomes

$$\mathcal{Z}_{\mathcal{G}}(\mathcal{G}) \propto \mathcal{S} \prod_{q \in Q_2} \mathcal{Z}_q, \tag{375}$$

showing that $(\mathcal{G}, \mathcal{S})$ is *not* a gauge code, but $(\mathcal{G}, \mathcal{S}')$ is, with

$$\mathcal{S}' := \mathcal{S} \prod_{q \in Q_2} \langle z_{1,q}, z_{2,q}\rangle. \tag{376}$$

The first equation in (295) becomes

$$\mathcal{Z}_{\mathcal{G}}(\mathcal{P}) = \mathcal{Z}_{\mathcal{G}}(\mathcal{G}) \prod_{q \in Q_1} \mathcal{Z}_q \mathcal{Z}^*_q, \tag{377}$$

showing that the encoded operators can be chosen to be, for $q \in Q_1$,

$$\begin{aligned} \bar X_q &:= z_{1,q}, & \bar Z_q &:= z^*_{1,q}, \\ \bar X_{\alpha+q} &:= z_{2,q}, & \bar Z_{\alpha+q} &:= z^*_{2,q}, \end{aligned} \tag{378}$$

That is, there are 2α encoded qubits.

Instead of making $\mathcal{S}$ non-local, we can take as the gauge group

$$\mathcal{G}' := \mathcal{G} \prod_{q\in Q_2} \langle z^*_{1,q}, z^*_{2,q}\rangle, \tag{379}$$

and then $(\mathcal{G}', \mathcal{S})$ is a gauge code too. In practice both choices are very similar, since the encoded operators are actually the same. The only difference appears in error correction, but for large systems the value of the extra stabilizers becomes too random and is thus of no use.

XI. CONCLUSIONS AND OUTLOOK

Under the sole assumption of translational invariance, we have provided a detailed study of the structure of two-dimensional TSCs. This structure can always be understood in terms of string operators that carry a topological charge, allowing to extend the insights from well known codes such as the toric code. Codes can be classified in terms of their topological charges, which are invariant under local transformations. In the case of subspace codes, two codes that have equivalent topological charges can be related by a local transformation. It remains open whether or not chiral subspace codes can be constructed.

We finish with a discussion of some natural extensions of the present work.

Boundaries — The same ideas that we have used to classify TSCs can be applied to classify boundaries between them. To model them, we can take the right and left half-plane to correspond to two possibly different TSSGs, allowing for arbitrary but translationally symmetric —along the axis direction— gauge and stabilizer generators in the axis between them. Of course, we have to impose the topological condition on $\mathcal{S}$, which still makes sense because we do not want to have localized degrees of freedom on small portions of the boundary.

Clearly there exist natural morphisms from the charge groups of the two TSSGs to the charge group of this boundary TSSG. These turn out to be enough to label all the charges —because no global constraints, and thus no charge, can be confined to the axis—. For charges not confined to either half-plane, the values of κ and θ on each side must agree. This is in particular true for the trivial charge. Thus, those charges that form the kernel of each of the two natural morphisms must be bosons with trivial mutual interactions. Such charges are especially relevant because they 'disolve' in the boundary, giving rise to logical string operators with endpoints on the boundary [11].

General codes — Although one could consider generalizations of these results to, say, stabilizer codes over qudits, it would be more interesting to investigate the structure of general two-dimensional topological quantum error correcting codes, defined in terms of LTI sets of commuting projectors. The expected result is that such codes would be describable in terms of anyon models with additional structure.

Higher dimensions — Even in the case of the relatively simple stabilizer formalism, the general structure of (translationally invariant) topological codes in three dimensions and above turns out to be quite rich. Already in the three-dimensional case there exist examples that do not fit on the standard homological picture [12, 14]. This is of great interesting due to the possible thermal stability [32] of the corresponding quantum memories, protected by the local Hamiltonian.

Acknowlegments— I would like to acknowledge useful discussions with Sergey Bravyi and Daniel Gottesman. I thank especially Guillaume Duclos-Cianci and David Poulin for many discussions and comments on several versions of the paper. This work was supported with research grants QUITEMAD S2009-ESP- 1594, FIS2009-10061 and UCM-BS/910758.

---

[1] P. Shor, Phys. Rev. A **52**, 2493 (1995).
[2] A. Steane, Phys. Rev. Lett. **77**, 793 (1996).
[3] E. Knill and R. Laflamme, Phys. Rev. A **55**, 900 (1997).
[4] C. Bennett, D. DiVincenzo, J. Smolin, and W. Wootters, Phys. Rev. A **54**, 3824 (1996).
[5] D. Gottesman, Phys. Rev. A **54**, 1862 (1996).
[6] A. Calderbank, E. Rains, P. Shor, and N. Sloane, Phys. Rev. Lett. **78**, 405 (1997).
[7] A. Kitaev, Ann. Phys. **303**, 2 (2003).
[8] E. Dennis, A. Kitaev, A. Landahl, and J. Preskill, J. Math. Phys. **43**, 4452 (2002).
[9] H. Bombin and M. A. Martin-Delgado, Phys. Rev. Lett. **97**, 180501 (2006).
[10] H. Bombin and M. A. Martin-Delgado, Phys. Rev. Lett. **98**, 160502 (2007).
[11] H. Bombin, Phys. Rev. A **81**, 032301 (2010).
[12] S. Bravyi, B. Leemhuis, and B. Terhal, Ann. of Phys. (2010).
[13] M. Suchara, S. Bravyi, and B. Terhal, J. of Phys. A: Math. and Theor. **44**, 155301 (2011).
[14] J. Haah, arXiv:1101.1962 (2011).
[15] D. Kribs, R. Laflamme, and D. Poulin, Phys. Rev. Lett. **94**, 180501 (2005).
[16] D. Poulin, Phys. Rev. Lett. **95**, 230504 (2005).
[17] D. Bacon, Phys. Rev. A **73**, 12340 (2006).
[18] C. Wang, J. Harrington, and J. Preskill, Ann. of Phys. **303**, 31 (2003).
[19] H. Katzgraber, H. Bombin, and M. Martin-Delgado, Phys. Rev. Lett. **103**, 090501 (2009).
[20] H. Katzgraber, H. Bombin, R. Andrist, and M. Martin-

Delgado, Phys. Rev. A (2010) **81**, 012319 (2010).

[21] R. Andrist, H. Katzgraber, H. Bombin, and M. Martin-Delgado, arXiv:1005.0777 (2010).

[22] T. Stace and S. Barrett, Phys. Rev. A **81**, 22317 (2010).

[23] B. Yoshida, Ann. of Phys. (2010).

[24] M. Freedman and D. Meyer, Found. of Comp. Math. **1**, 325 (2001).

[25] H. Bombin, Phys. Rev. Lett. **105**, 30403 (2010).

[26] R. Raussendorf, J. Harrington, and K. Goyal, New J. Phys. **9**, 199 (2007).

[27] H. Bombin and M. A. Martin-Delgado, J. Phys. A: Math. and Theor. **42**, 095302 (2009).

[28] G. Duclos-Cianci and D. Poulin, Phys. Rev. Lett. **104**, 50504 (2010).

[29] S. Bravyi and B. Terhal, New J. Phys. **11**, 043029 (2009).

[30] S. Bravyi, D. Poulin, and B. Terhal, Phys. Rev. Lett. **104**, 50503 (2010).

[31] S. Bravyi, Phys. Rev. A **83**, 012320 (2011).

[32] S. Bravyi and J. Haah, arXiv:1105.4159 (2011).

[33] H. Bombin, G. Duclos-Cianci, and D. Poulin, Arxiv preprint arXiv:1103.4606 (2011).

[34] M. Levin and X.-G. Wen, Phys. Rev. B **67**, 245316 (2003).